

09/24/99  
Jc583 U.S. PTO

Practitioner's Docket No. 48551

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

Jc678 U.S. PTO  
09/406269  
09/24/99

NEW APPLICATION TRANSMITTAL

Transmitted herewith for filing is the patent application of

Inventor(s): Jin-An Jiao, Lawrence K. Luepschen, Esperanza L. Nieves,  
Hing C. Wong and Dean P. Taylor

**WARNING:** 37 C.F.R. § 1.41(a)(1) points out:

"(a) A patent is applied for in the name or names of the actual inventor or inventors.

"(1) The inventorship of a nonprovisional application is that inventorship set forth in the oath or declaration as prescribed by § 1.63, except as provided for in § 1.53(d)(4) and § 1.63(d). If an oath or declaration as prescribed by § 1.63 is not filed during the pendency of a nonprovisional application, the inventorship is that inventorship set forth in the application papers filed pursuant to § 1.53(b), unless a petition under this paragraph accompanied by the fee set forth in § 1.17(i) is filed supplying or changing the name or names of the inventor or inventors."

For (title):

PHARMACEUTICALLY ACTIVE COMPOUNDS AND METHODS OF USE THEREOF

**CERTIFICATION UNDER 37 C.F.R. § 1.10\***

(Express Mail label number is **mandatory**.)

(Express Mail certification is **optional**.)

I hereby certify that this New Application Transmittal and the documents referred to as attached therein are being deposited with the United States Postal Service on this date September 24, 1999, in an envelope as "Express Mail Post Office to Addressee," mailing Label Number EI977542427US, addressed to the: Assistant Commissioner for Patents, Washington, D.C. 20231.

Deanna M. Landry

(type or print name of person mailing paper)

Deanna Landry  
Signature of person mailing paper

**WARNING:** Certificate of mailing (first class) or facsimile transmission procedures of 37 C.F.R. § 1.8 cannot be used to obtain a date of mailing or transmission for this correspondence.

**\*WARNING:** Each paper or fee filed by "Express Mail" **must** have the number of the "Express Mail" mailing label placed thereon prior to mailing. 37 C.F.R. § 1.10(b).

"Since the filing of correspondence under § 1.10 without the Express Mail mailing label thereon is an oversight that can be avoided by the exercise of reasonable care, requests for waiver of this requirement will **not** be granted on petition." Notice of Oct. 24, 1996, 60 Fed. Reg. 56,439, at 56,442.

(New Application Transmittal [4-1]—page 1 of 11)

## 1. Type of Application

This new application is for a(n)

(check one applicable item below)

- ☒ Original (nonprovisional)  
☐ Design  
☐ Plant

**WARNING:** Do not use this transmittal for a completion in the U.S. of an International Application under 35 U.S.C. § 371(c)(4), unless the International Application is being filed as a divisional, continuation or continuation-in-part application.

**WARNING:** Do not use this transmittal for the filing of a provisional application.

**NOTE:** If one of the following 3 items apply, then complete and attach **ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF A PRIOR U.S. APPLICATION CLAIMED** and a **NOTIFICATION IN PARENT APPLICATION OF THE FILING OF THIS CONTINUATION APPLICATION**.

- ☐ Divisional.  
☐ Continuation.  
☐ Continuation-in-part (C-I-P).

## 2. Benefit of Prior U.S. Application(s) (35 U.S.C. §§ 119(e), 120, or 121)

**NOTE:** A nonprovisional application may claim an invention disclosed in one or more prior filed copending nonprovisional applications or copending international applications designating the United States of America. In order for a nonprovisional application to claim the benefit of a prior filed copending nonprovisional application or copending international application designating the United States of America, each prior application must name as an inventor at least one inventor named in the later filed nonprovisional application and disclose the named inventor's invention claimed in at least one claim of the later filed nonprovisional application in the manner provided by the first paragraph of 35 U.S.C. § 112. Each prior application must also be:

(i) An international application entitled to a filing date in accordance with PCT Article 11 and designating the United States of America; or

(ii) Complete as set forth in § 1.51(b); or

(iii) Entitled to a filing date as set forth in § 1.53(b) or § 1.53(d) and include the basic filing fee set forth in § 1.16; or

(iv) Entitled to a filing date as set forth in § 1.53(b) and have paid therein the processing and retention fee set forth in § 1.21(f) within the time period set forth in § 1.53(f).

37 C.F.R. § 1.78(a)(1).

**NOTE:** If the new application being transmitted is a divisional, continuation or a continuation-in-part of a parent case, or where the parent case is an International Application which designated the U.S., or benefit of a prior provisional application is claimed, then check the following item and complete and attach **ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED**.

**WARNING:** If an application claims the benefit of the filing date of an earlier filed application under 35 U.S.C. §§ 120, 121 or 365(c), the 20-year term of that application will be based upon the filing date of the earliest U.S. application that the application makes reference to under 35 U.S.C. §§ 120, 121 or 365(c). (35 U.S.C. § 154(a)(2) does not take into account, for the determination of the patent term, any application on which priority is claimed under 35 U.S.C. §§ 119, 365(a) or 365(b).) For a c-i-p application, applicant should review whether any claim in the patent that will issue is supported by an earlier application and, if not, the applicant should consider canceling the reference to the earlier filed application. The term of a patent is not based on a claim-by-claim approach. See Notice of April 14, 1995, 60 Fed. Reg. 20,195, at 20,205.

(New Application Transmittal [4-1]—page 2 of 11)

**WARNING:** When the last day of pendency of a provisional application falls on a Saturday, Sunday, or Federal holiday within the District of Columbia, any nonprovisional application claiming benefit of the provisional application **must** be filed prior to the Saturday, Sunday, or Federal holiday within the District of Columbia. See 37 C.F.R. § 1.78(a)(3).

- ☒ The new application being transmitted claims the benefit of prior U.S. application(s). Enclosed are ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED.

### 3. Papers Enclosed

- A. Required for filing date under 37 C.F.R. § 1.53(b) (Regular) or 37 C.F.R. § 1.153 (Design) Application

36 Pages of specification

17 Pages of claims

10 Sheets of drawing

**WARNING:** **DO NOT** submit original drawings. A high quality copy of the drawings should be supplied when filing a patent application. The drawings that are submitted to the Office must be on strong, white, smooth, and non-shiny paper and meet the standards according to § 1.84. If corrections to the drawings are necessary, they should be made to the original drawing and a high-quality copy of the corrected original drawing then submitted to the Office. Only one copy is required or desired. For comments on proposed then-new 37 C.F.R. § 1.84, see Notice of March 9, 1988 (1990 O.G. 57-62).

**NOTE:** "Identifying indicia, if provided, should include the application number or the title of the invention, inventor's name, docket number (if any), and the name and telephone number of a person to call if the Office is unable to match the drawings to the proper application. This information should be placed on the back of each sheet of drawing a minimum distance of 1.5 cm. (5/8 inch) down from the top of the page . . ." 37 C.F.R. § 1.84(c)).

(complete the following, if applicable)

- ☐ The enclosed drawing(s) are photograph(s), and there is also attached a "PETITION TO ACCEPT PHOTOGRAPH(S) AS DRAWING(S)." 37 C.F.R. § 1.84(b).
- ☐ formal
- ☐ informal

### B. Other Papers Enclosed

       Pages of declaration and power of attorney

  1   Pages of abstract

       Other

### 4. Additional papers enclosed

- ☐ Amendment to claims
- ☐ Cancel in this applications claims \_\_\_\_\_ before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)
- ☐ Add the claims shown on the attached amendment. (Claims added have been numbered consecutively following the highest numbered original claims.)
- ☐ Preliminary Amendment
- ☐ Information Disclosure Statement (37 C.F.R. § 1.98)
- ☐ Form PTO-1449 (PTO/SB/08A and 08B)
- ☐ Citations

- ☐ Declaration of Biological Deposit
- ☐ Submission of "Sequence Listing," computer readable copy and/or amendment pertaining thereto for biotechnology invention containing nucleotide and/or amino acid sequence.
- ☐ Authorization of Attorney(s) to Accept and Follow Instructions from Representative
- ☐ Special Comments
- ☐ Other

#### 5. Declaration or oath (including power of attorney)

**NOTE:** A newly executed declaration is not required in a continuation or divisional application provided that the prior nonprovisional application contained a declaration as required, the application being filed is by all or fewer than all the inventors named in the prior application, there is no new matter in the application being filed, and a copy of the executed declaration filed in the prior application (showing the signature or an indication thereon that it was signed) is submitted. The copy must be accompanied by a statement requesting deletion of the names of person(s) who are not inventors of the application being filed. If the declaration in the prior application was filed under § 1.47, then a copy of that declaration must be filed accompanied by a copy of the decision granting § 1.47 status or, if a nonsigning person under § 1.47 has subsequently joined in a prior application, then a copy of the subsequently executed declaration must be filed. See 37 C.F.R. §§ 1.63(d)(1)–(3).

**NOTE:** A declaration filed to complete an application must be executed, identify the specification to which it is directed, identify each inventor by full name including family name and at least one given name, without abbreviation together with any other given name or initial, and the residence, post office address and country or citizenship of each inventor, and state whether the inventor is a sole or joint inventor. 37 C.F.R. § 1.63(a)(1)–(4).

- ☐ Enclosed
- Executed by

(check all applicable boxes)

- ☐ inventor(s).
- ☐ legal representative of inventor(s).  
37 C.F.R. §§ 1.42 or 1.43.
- ☐ joint inventor or person showing a proprietary interest on behalf of inventor who refused to sign or cannot be reached.
- ☐ This is the petition required by 37 C.F.R. § 1.47 and the statement required by 37 C.F.R. § 1.47 is also attached. See item 13 below for fee.

- ☒ Not Enclosed.

**NOTE:** Where the filing is a completion in the U.S. of an International Application or where the completion of the U.S. application contains subject matter in addition to the International Application, the application may be treated as a continuation or continuation-in-part, as the case may be, utilizing ADDED PAGE FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION CLAIMED.

- ☐ Application is made by a person authorized under 37 C.F.R. § 1.41(c) on behalf of all the above named inventor(s).

(The declaration or oath, along with the surcharge required by 37 C.F.R. § 1.16(e) can be filed subsequently).

- ☐ Showing that the filing is authorized.  
(not required unless called into question. 37 C.F.R. § 1.41(d))

## 6. Inventorship Statement

**WARNING:** If the named inventors are each not the inventors of all the claims an explanation, including the ownership of the various claims at the time the last claimed invention was made, should be submitted.

The inventorship for all the claims in this application are:

☐ The same.

or

☐ Not the same. An explanation, including the ownership of the various claims at the time the last claimed invention was made,

☐ is submitted.

☐ will be submitted.

## 7. Language

**NOTE:** An application including a signed oath or declaration may be filed in a language other than English. An English translation of the non-English language application and the processing fee of \$130.00 required by 37 C.F.R. § 1.17(k) is required to be filed with the application, or within such time as may be set by the Office. 37 C.F.R. § 1.52(d).

☒ English

☐ Non-English

☐ The attached translation includes a statement that the translation is accurate. 37 C.F.R. § 1.52(d).

## 8. Assignment

☒ An assignment of the invention to Sunol Molecular Corporation  
of Miramar, Florida

☐ is attached. A separate ☐ "COVER SHEET FOR ASSIGNMENT (DOCUMENT) ACCOMPANYING NEW PATENT APPLICATION" or ☐ FORM PTO 1595 is also attached.

☒ will follow.

**NOTE:** "If an assignment is submitted with a new application, send two separate letters—one for the application and one for the assignment." Notice of May 4, 1990 (1114 O.G. 77-78).

**WARNING:** A newly executed "CERTIFICATE UNDER 37 C.F.R. § 3.73(b)" must be filed when a continuation-in-part application is filed by an assignee. Notice of April 30, 1993, 1150 O.G. 62-64.

(New Application Transmittal [4-1]—page 5 of 11)

## 9. Certified Copy

Certified copy(ies) of application(s)

Country	Appln. No.	Filed
Country	Appln. No.	Filed
Country	Appln. No.	Filed

from which priority is claimed

☐ is (are) attached.

☐ will follow.

NOTE: The foreign application forming the basis for the claim for priority must be referred to in the oath or declaration. 37 C.F.R. § 1.55(a) and 1.63.

NOTE: This item is for any foreign priority for which the application being filed directly relates. If any parent U.S. application or International Application from which this application claims benefit under 35 U.S.C. § 120 is itself entitled to priority from a prior foreign application, then complete item 18 on the ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED.

## 10. Fee Calculation (37 C.F.R. § 1.16)

A. ☒ Regular application

CLAIMS AS FILED			
Number filed	Number Extra	Rate	Basic Fee 37 C.F.R. 1.16(a) \$760.00
Total Claims (37 C.F.R. § 1.16(c))	— 20 =	×	\$ 18.00
Independent Claims (37 C.F.R. § 1.16(b))	— 3 =	×	\$ 78.00
Multiple dependent claim(s), if any (37 C.F.R. § 1.16(d))		+	\$260.00

☐ Amendment cancelling extra claims is enclosed.

☐ Amendment deleting multiple-dependencies is enclosed.

☐ Fee for extra claims is not being paid at this time.

NOTE: If the fees for extra claims are not paid on filing they must be paid or the claims cancelled by amendment, prior to the expiration of the time period set for response by the Patent and Trademark Office in any notice of fee deficiency. 37 C.F.R. § 1.16(d).

Filing Fee Calculation \$ \_\_\_\_\_

B. ☐ Design application  
(\$310.00—37 C.F.R. § 1.16(f))

Filing Fee Calculation \$ \_\_\_\_\_

C. ☐ Plant application  
(\$480.00—37 C.F.R. § 1.16(g))

Filing fee calculation \$ \_\_\_\_\_

11. Small Entity Statement(s)

- ☐ Statement(s) that this is a filing by a small entity under 37 C.F.R. § 1.9 and 1.27 is (are) attached.

**WARNING:** "Status as a small entity must be specifically established in each application or patent in which the status is available and desired. Status as a small entity in one application or patent does not affect any other application or patent, including applications or patents which are directly or indirectly dependent upon the application or patent in which the status has been established. The refiling of an application under § 1.53 as a continuation, division, or continuation-in-part (including a continued prosecution application under § 1.53(d)), or the filing of a reissue application requires a new determination as to continued entitlement to small entity status for the continuing or reissue application. A nonprovisional application claiming benefit under 35 U.S.C. § 119(e), 120, 121, or 365(c) of a prior application, or a reissue application may rely on a statement filed in the prior application or in the patent if the nonprovisional application or the reissue application includes a reference to the statement in the prior application or in the patent or includes a copy of the statement in the prior application or in the patent and status as a small entity is still proper and desired. The payment of the small entity basic statutory filing fee will be treated as such a reference for purposes of this section." 37 C.F.R. § 1.28(a)(2).

**WARNING:** "Small entity status must not be established when the person or persons signing the . . . statement can **unequivocally** make the required self-certification." M.P.E.P., § 509.03, 6th ed., rev. 2, July 1996 (emphasis added).

(complete the following, if applicable)

- ☐ Status as a small entity was claimed in prior application \_\_\_\_\_ / \_\_\_\_\_, filed on \_\_\_\_\_, from which benefit is being claimed for this application under:

35 U.S.C. § ☐ 119(e),  
☐ 120,  
☐ 121,  
☐ 365(c),

and which status as a small entity is still proper and desired.

- ☐ A copy of the statement in the prior application is included.

Filing Fee Calculation (50% of A, B or C above)

\$ \_\_\_\_\_

**NOTE:** Any excess of the full fee paid will be refunded if small entity status is established and a refund request are filed within 2 months of the date of timely payment of a full fee. The two-month period is not extendable under § 1.136. 37 C.F.R. § 1.28(a).

12. Request for International-Type Search (37 C.F.R. § 1.104(d))

(complete, if applicable)

- ☐ Please prepare an international-type search report for this application at the time when national examination on the merits takes place.

**13. Fee Payment Being Made at This Time**

☒ Not Enclosed

☒ No filing fee is to be paid at this time.

*(This and the surcharge required by 37 C.F.R. § 1.16(e) can be paid subsequently.)*

☐ Enclosed

☐ Filing fee \$ \_\_\_\_\_

☐ Recording assignment  
(\$40.00; 37 C.F.R. § 1.21(h))  
(See attached "COVER SHEET FOR  
ASSIGNMENT ACCOMPANYING NEW  
APPLICATION".) \$ \_\_\_\_\_

☐ Petition fee for filing by other than all the  
inventors or person on behalf of the inventor  
where inventor refused to sign or cannot be  
reached  
(\$130.00; 37 C.F.R. §§ 1.47 and 1.17(l)) \$ \_\_\_\_\_

☐ For processing an application with a  
specification in  
a non-English language  
(\$130.00; 37 C.F.R. §§ 1.52(d) and 1.17(k)) \$ \_\_\_\_\_

☐ Processing and retention fee  
(\$130.00; 37 C.F.R. §§ 1.53(d) and 1.21(l)) \$ \_\_\_\_\_

☐ Fee for international-type search report  
(\$40.00; 37 C.F.R. § 1.21(e)) \$ \_\_\_\_\_

**NOTE:** 37 C.F.R. § 1.21(l) establishes a fee for processing and retaining any application that is abandoned for failing to complete the application pursuant to 37 C.F.R. § 1.53(f) and this, as well as the changes to 37 C.F.R. §§ 1.53 and 1.78(a)(1), indicate that in order to obtain the benefit of a prior U.S. application, either the basic filing fee must be paid, or the processing and retention fee of § 1.21(l) must be paid, within 1 year from notification under § 53(f).

Total fees enclosed \$ \_\_\_\_\_

**14. Method of Payment of Fees**

☐ Check in the amount of \$ \_\_\_\_\_

☐ Charge Account No. \_\_\_\_\_ in the amount of  
\$ \_\_\_\_\_.

A duplicate of this transmittal is attached.

**NOTE:** Fees should be itemized in such a manner that it is clear for which purpose the fees are paid. 37 C.F.R. § 1.22(b).



## 15. Authorization to Charge Additional Fees

**WARNING:** If no fees are to be paid on filing, the following items should not be completed.

**WARNING:** Accurately count claims, especially multiple dependent claims, to avoid unexpected high charges, if extra claim charges are authorized.

- ☐ The Commissioner is hereby authorized to charge the following additional fees by this paper and during the entire pendency of this application to Account No. \_\_\_\_\_:

- ☐ 37 C.F.R. § 1.16(a), (f) or (g) (filing fees)  
☐ 37 C.F.R. § 1.16(b), (c) and (d) (presentation of extra claims)

**NOTE:** Because additional fees for excess or multiple dependent claims not paid on filing or on later presentation must only be paid or these claims cancelled by amendment prior to the expiration of the time period set for response by the PTO in any notice of fee deficiency (37 C.F.R. § 1.16(d)), it might be best not to authorize the PTO to charge additional claim fees, except possibly when dealing with amendments after final action.

- ☐ 37 C.F.R. § 1.16(e) (surcharge for filing the basic filing fee and/or declaration on a date later than the filing date of the application)  
☐ 37 C.F.R. § 1.17(a)(1)–(5) (extension fees pursuant to § 1.136(a)).  
☐ 37 C.F.R. § 1.17 (application processing fees)

**NOTE:** “. . . A written request may be submitted in an application that is an authorization to treat any concurrent or future reply, requiring a petition for an extension of time under this paragraph for its timely submission, as incorporating a petition for extension of time for the appropriate length of time. An authorization to charge all required fees, fees under § 1.17, or all required extension of time fees will be treated as a constructive petition for an extension of time in any concurrent or future reply requiring a petition for an extension of time under this paragraph for its timely submission. Submission of the fee set forth in § 1.17(a) will also be treated as a constructive petition for an extension of time in any concurrent reply requiring a petition for an extension of time under this paragraph for its timely submission.” 37 C.F.R. § 1.136(a)(3).

- ☐ 37 C.F.R. § 1.18 (issue fee at or before mailing of Notice of Allowance, pursuant to 37 C.F.R. § 1.311(b))

**NOTE:** Where an authorization to charge the issue fee to a deposit account has been filed before the mailing of a Notice of Allowance, the issue fee will be automatically charged to the deposit account at the time of mailing the notice of allowance. 37 C.F.R. § 1.311(b).

**NOTE:** 37 C.F.R. § 1.28(b) requires “Notification of any change in status resulting in loss of entitlement to small entity status must be filed in the application . . . prior to paying, or at the time of paying, . . . the issue fee. . . .” From the wording of 37 C.F.R. § 1.28(b), (a) notification of change of status must be made even if the fee is paid as “other than a small entity” and (b) no notification is required if the change is to another small entity.

**16. Instructions as to Overpayment**


**NOTE:** "... Amounts of twenty-five dollars or less will not be returned unless specifically requested within a reasonable time, nor will the payer be notified of such amounts; amounts over twenty-five dollars may be returned by check or, if requested, by credit to a deposit account." 37 C.F.R. § 1.26(a).

- ☐ Credit Account No. \_\_\_\_\_
- ☐ Refund

Reg. No. 33,860

Tel. No. (617) 523-3400

Customer No.



SIGNATURE OF PRACTITIONER

Peter F. Corless

(type or print name of attorney)

130 Water Street

P.O. Address

Boston, MA 02109

☒ **Incorporation by reference of added pages**

*(check the following item if the application in this transmittal claims the benefit of prior U.S. application(s) (including an international application entering the U.S. stage as a continuation, divisional or C-I-P application) and complete and attach the ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED)*

- ☒ Plus Added Pages for New Application Transmittal Where Benefit of Prior U.S. Application(s) Claimed

Number of pages added 5

- ☐ Plus Added Pages for Papers Referred to in Item 4 Above

Number of pages added \_\_\_\_\_

- ☐ Plus added pages deleting names of inventor(s) named in prior application(s) who is/are no longer inventor(s) of the subject matter claimed in this application.

Number of pages added \_\_\_\_\_

- ☐ Plus "Assignment Cover Letter Accompanying New Application"

Number of pages added \_\_\_\_\_

☐ **Statement Where No Further Pages Added**

*(if no further pages form a part of this Transmittal, then end this Transmittal with this page and check the following item)*

- ☐ This transmittal ends with this page.

ADDED PAGES FOR APPLICATION TRANSMITTAL WHERE BENEFIT OF  
PRIOR U.S. APPLICATION(S) CLAIMED

NOTE: See 37 CFR 1.78.

**17. Relate Back**

**WARNING:** If an application claims the benefit of the filing date of an earlier filed application under 35 U.S.C. 120, 121 or 365(c), the 20-year term of that application will be based upon the filing date of the earliest U.S. application that the application makes reference to under 35 U.S.C. 120, 121 or 365(c). (35 U.S.C. 154(a)(2) does not take into account, for the determination of the patent term, any application on which priority is claimed under 35 U.S.C. 119, 365(a) or 365(b).) For a c-i-p application, applicant should review whether any claim in the patent that will issue is supported by an earlier application and, if not, the applicant should consider canceling the reference to the earlier filed application. The term of a patent is not based on a claim-by-claim approach. See Notice of April 14, 1995, 60 Fed. Reg. 20,195, at 20,205.

(complete the following, if applicable)

☒ Amend the specification by inserting, before the first line, the following sentence:**A. 35 U.S.C. 119(e)**

NOTE: "Any nonprovisional application claiming the benefit of one or more prior filed copending provisional applications must contain or be amended to contain in the first sentence of the specification following the title a reference to each such prior provisional application, identifying it as a provisional application, and including the provisional application number (consisting of series code and serial number)." 37 C.F.R. § 1.78(a)(4).

☒ "This application claims the benefit of U.S. Provisional Application(s) No(s).:**APPLICATION NO(S).:****FILING DATE**

<u>60</u> / <u>101,887</u>	<u>09/25/98</u> "
<u>      </u> / <u>                    </u>	<u>                    </u> "
<u>      </u> / <u>                    </u>	<u>                    </u> "

**B. 35 U.S.C. 120, 121 and 365(c)**

NOTE: "Except for a continued prosecution application filed under § 1.53(d), any nonprovisional application claiming the benefit of one or more prior filed copending nonprovisional applications or international applications designating the United States of America must contain or be amended to contain in the first sentence of the specification following the title a reference to each such prior application, identifying it by application number (consisting of the series code and serial number) or international application number and international filing date and indicating the relationship of the applications. Cross-references to other related applications may be made when appropriate." (See § 1.14(a)). 37 C.F.R. § 1.78(a)(2).

- ☐ "This application is a  
☐ continuation  
☐ continuation-in-part  
☐ divisional

of copending application(s)

- ☐ application number 0 / \_\_\_\_\_ filed on \_\_\_\_\_"  
☐ International Application \_\_\_\_\_ filed on \_\_\_\_\_  
\_\_\_\_\_ and which designated the U.S."

NOTE: The proper reference to a prior filed PCT application that entered the U.S. national phase is the U.S. serial number and the filing date of the PCT application that designated the U.S.

NOTE: (1) Where the application being transmitted adds subject matter to the International Application, then the filing can be as a continuation-in-part or (2) if it is desired to do so for other reasons then the filing can be as a continuation.

NOTE: The deadline for entering the national phase in the U.S. for an international application was clarified in the Notice of April 28, 1987 (1079 O.G. 32 to 46) as follows:

"The Patent and Trademark Office considers the International application to be pending until the 22nd month from the priority date if the United States has been designated and no Demand for International Preliminary Examination has been filed prior to the expiration of the 19th month from the priority date and until the 32nd month from the priority date if a Demand for International Preliminary Examination which elected the United States of America has been filed prior to the expiration of the 19th month from the priority date, provided that a copy of the international application has been communicated to the Patent and Trademark Office within the 20 or 30 month period respectively. If a copy of the international application has not been communicated to the Patent and Trademark Office within the 20 or 30 month period respectively, the international application becomes abandoned as to the United States 20 or 30 months from the priority date respectively. These periods have been placed in the rules as paragraph (h) of § 1.494 and paragraph (i) of § 1.495. A continuing application under 35 U.S.C. 365(c) and 120 may be filed anytime during the pendency of the international application."

- ☐ "The nonprovisional application designated above, namely application \_\_\_\_\_ / \_\_\_\_\_, filed \_\_\_\_\_, claims the benefit of U.S. Provisional Application(s) No(s).:

**APPLICATION NO(S):**

**FILING DATE**

_____ / _____	_____ "
_____ / _____	_____ "
_____ / _____	_____ "

- ☐ Where more than one reference is made above, please combine all references into one sentence.

### 18. Relate Back—35 U.S.C. 119 Priority Claim for Prior Application

The prior U.S. application(s), including any prior International Application designating the U.S., identified above in item 17B, in turn itself claim(s) foreign priority(ies) as follows:

Country	Appin. no.	Filed on
---------	------------	----------

The certified copy(ies) has (have)

- ☐ been filed on \_\_\_\_\_, in prior application 0 / \_\_\_\_\_, which was filed on \_\_\_\_\_.
- ☐ is (are) attached.

**WARNING:** *The certified copy of the priority application that may have been communicated to the PTO by the International Bureau may not be relied on without any need to file a certified copy of the priority application in the continuing application. This is so because the certified copy of the priority application communicated by the International Bureau is placed in a folder and is not assigned a U.S. serial number unless the national stage is entered. Such folders are disposed of if the national stage is not entered. Therefore, such certified copies may not be available if needed later in the prosecution of a continuing application. An alternative would be to physically remove the priority documents from the folders and transfer them to the continuing application. The resources required to request transfer, retrieve the folders, make suitable record notations, transfer the certified copies, enter and make a record of such copies in the Continuing Application are substantial. Accordingly, the priority documents in folders of international applications that have not entered the national stage may not be relied on. Notice of April 28, 1987 (1079 O.G. 32 to 46).*

### 19. Maintenance of Copendency of Prior Application

**NOTE:** *The PTO finds it useful if a copy of the petition filed in the prior application extending the term for response is filed with the papers constituting the filing of the continuation application. Notice of November 5, 1985 (1060 O.G. 27).*

- A.** ☐ Extension of time in prior application

*(This item must be completed and the papers filed in the prior application, if the period set in the prior application has run.)*

- ☐ A petition, fee and response extends the term in the pending **prior** application until \_\_\_\_\_.
- ☐ A **copy** of the petition filed in prior application is attached.

- B.** ☐ Conditional Petition for Extension of Time in Prior Application

*(complete this item, if previous item not applicable)*

- ☐ A conditional petition for extension of time is being filed in the pending **prior** application.
- ☐ A **copy** of the conditional petition filed in the prior application is attached.

**20. Further Inventorship Statement Where Benefit of Prior Application(s) Claimed**

*(complete applicable item (a), (b) and/or (c) below)*

- (a) ☐ This application discloses and claims only subject matter disclosed in the prior application whose particulars are set out above and the inventor(s) in this application are
- ☐ the same.
  - ☐ less than those named in the prior application. It is requested that the following inventor(s) identified for the prior application be deleted:

\_\_\_\_\_  
*(type name(s) of inventor(s) to be deleted)*

- (b) ☐ This application discloses and claims additional disclosure by amendment and a new declaration or oath is being filed. With respect to the prior application, the inventor(s) in this application are
- ☐ the same.
  - ☐ the following additional inventor(s) have been added:

\_\_\_\_\_  
*(type name(s) of inventor(s) to be added)*

- (c) The inventorship for all the claims in this application are
- ☐ the same.
  - ☐ not the same. An explanation, including the ownership of the various claims at the time the last claimed invention was made
    - ☐ is submitted.
    - ☐ will be submitted.

**21. Abandonment of Prior Application (if applicable)**

- ☐ Please abandon the prior application at a time while the prior application is pending, or when the petition for extension of time or to revive in that application is granted, and when this application is granted a filing date, so as to make this application copending with said prior application.

*NOTE: According to the Notice of May 13, 1983 (103, TMOG 6-7), the filing of a continuation or continuation-in-part application is a proper response with respect to a petition for extension of time or a petition to revive and should include the express abandonment of the prior application conditioned upon the granting of the petition and the granting of a filing date to the continuing application.*

**22. Petition for Suspension of Prosecution for the Time Necessary to File an Amendment**

**WARNING:** *"The claims of a new application may be finally rejected in the first Office action in those situations where (1) the new application is a continuing application of, or a substitute for, an earlier application, and (2) all the claims of the new application (a) are drawn to the same invention claimed in the earlier application, and (b) would have been properly finally rejected on the grounds of art of record in the next Office action if they had been entered in the earlier application." MPEP, § 706.07(b), 6th ed., rev.2.*

*NOTE: Where it is possible that the claims on file will give rise to a first action final for this continuation application and for some reason an amendment cannot be filed promptly (e.g., experimental data is being gathered) it may be desirable to file a petition for suspension of prosecution for the time necessary.*

*(check the next item, if applicable)*

- ☐ There is provided herewith a Petition To Suspend Prosecution for the Time Necessary to File An Amendment (New Application Filed Concurrently)

**23. Small Entity (37 CFR § 1.28(a))**

- ☐ Applicant has established small entity status by the filing of a statement in parent application /\_\_\_\_\_ on \_\_\_\_\_.
- ☐ A copy of the statement previously filed is included.

**WARNING:** See 37 CFR § 1.28(a).

**24. NOTIFICATION IN PARENT APPLICATION OF THIS FILING**

- ☐ A notification of the filing of this  
*(check one of the following)*
- ☐ continuation
  - ☐ continuation-in-part
  - ☐ divisional

is being filed in the parent application, from which this application claims priority under 35 U.S.C. § 120.



PHARMACEUTICALLY ACTIVE COMPOUNDS  
AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATION

The present application claims the benefit of U. S. Provisional Application No. 60/101,887 filed on September 25, 1998, the disclosure of which is hereby incorporated by reference.

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to pharmaceutically active compounds and more particularly to pharmaceutical compositions that utilize or comprise one or more of such compounds. Preferred compounds are especially useful for the treatment or prophylaxis of undesired thrombosis. Also included are methods for treating thrombosis. The invention has a wide spectrum of applications including use in screening candidate compounds for the treatment or prophylaxis of thrombosis.

15 2. Background

Blood clotting assists hemostasis by minimizing blood loss. Generally, blood clotting is initiated by vessel damage and requires platelet aggregation, coagulation factors and inhibition of fibrinolysis. The coagulation factors act through a cascade that relates the vessel damage to formation of a blood clot (see generally L. Stryer, *Biochemistry*, 3rd Ed, W.H. Freeman Co., New York; and A.G. Gilman et al., *The Pharmacological Basis of Therapeutics*, 8th Edition, McGraw Hill Inc., New York, pp. 1311-1331).

Tissue factor (TF), an integral membrane protein of 263 amino acids, is responsible for initiating the coagulation protease cascade. Vascular damage exposes blood to tissue factor expressed on subendothelial cell surfaces, leading to the formation of a calcium-dependent, high-affinity complex with the plasma factor VII

25

(FVII) or activated factor VII (FVIIa). Binding to TF promotes rapid proteolytic cleavage of the zymogen FVII to the active serine protease FVIIa by a number of proteases such as factor Xa, or thrombin. TF also functions as an essential cofactor for FVIIa by dramatically enhancing the catalytic efficiency of FVIIa for its protein substrates factors IX and X. TF/VIIa complex activates factors IX (FIX) and X (FX) via limited proteolysis, ultimately leading to thrombin generation and fibrin deposition. Under pathological conditions such as atherosclerosis or following invasive surgical procedures such as microvascular graft, angioplasty, deployment of an implanted device (e.g., a stent, catheter or arteriovenous shunt), or endarterectomy, TF-initiated coagulation can lead to thrombotic disorders that can result e.g. in heart attack, stroke, unstable angina or other coronary disorder.

Thrombosis also may accompany various thromboembolic disorders and coagulopathies such as a pulmonary embolism (e.g., atrial fibrillation with embolization, deep vein thrombosis, *etc.*) and disseminated intravascular coagulation, respectively. Manipulation of body fluids can also result in an undesirable thrombus, particularly in blood transfusions or fluid sampling, as well as procedures involving extracorporeal circulation (e.g., cardiopulmonary bypass surgery) and dialysis.

Certain anti-coagulants have been used to alleviate or avoid blood clots associated with thrombosis. Blood clotting often can be minimized or eliminated by administering a suitable anti-coagulant or mixture thereof, including one or more of a coumarin derivative (e.g., warfarin and dicumarol) or a charged polymer (e.g., heparin, hirudin or hirulog). See e.g., Gilman et al., *supra*, R.J. Beigering et al., *Ann. Hematol.*, **72**:177 (1996); J.D. Willerson, *Circulation*, **94**:866 (1996).

Certain antibodies with anti-platelet activity have also been used to alleviate various thromboses. For example, ReoPro™ is a therapeutic antibody that is routinely administered to alleviate various thromboembolic disorders such as those arising from angioplasty, myocardial infarction, unstable angina and coronary artery stenoses. Additionally, ReoPro™ can be used as a prophylactic to reduce the risk of myocardial

infarction and angina (J.T. Willerson, *Circulation*, **94**:866 (1996); M.L. Simmons et al., *Circulation*, **89**:596 (1994)).

However, use of prior anti-coagulants is often associated with side effects such as hemorrhaging, re-occlusion, "white-clot" syndrome, irritation, birth defects, thrombocytopenia and hepatic dysfunction. Long-term administration of anti-coagulants can particularly increase risk of life-threatening illness (see e.g., Gilman et al., *supra*).

Protein-based agents are potentially safer, but generally are limited to treatment of acute conditions and often are restricted to parenteral administration. Such agents are considered less suitable for long-term therapies for chronic diseases (such as atherosclerosis, a major cause of heart attack) due to the increased probability of immune response to a protein therapeutic, relatively high production cost and/or limited oral bioavailability.

It would thus be desirable to have new anti-coagulant agents. It would be particularly desirable to have new anti-coagulant agents that could be administered over a relatively long period to treat chronic conditions such as atherosclerosis.

#### SUMMARY OF THE INVENTION

We have now discovered pharmaceutically active compounds and compositions that are useful to treat or prevent undesired thrombosis. Preferred compounds of the invention are tissue factor (TF) antagonists that preferably specifically block human factor X and IX activation catalyzed by human tissue factor/factor VIIa complex. Also discovered are methods for treating or preventing thrombosis that use the compounds and compositions disclosed herein.

More particular methods of this invention include administering a therapeutically effective amount of at least one compound or composition of this invention. The compound or composition is typically given to a mammal in need of such treatment such as a human patient who is susceptible to, suffering from, or

recovering from undesired thrombosis, or mammal that is suffering from, recovering from or susceptible to other disease or disorder impacted by tissue factor such as a cardiovascular disease, cell proliferation disorder, post-operative complication, or an immune disorder. Preferred compounds and compositions may also be used to treat or prevent recognized disorders impacted by various thromboses such as those particular disorders disclosed herein.

The invention also includes methods for blocking or inhibiting tissue factor-dependent activation of factor X and/or factor IX. These methods in general include contacting tissue factor with a TF blocking compound to thereby inhibit formation of a functional complex of factor X or factor IX with tissue factor or TF/VIIa. Preferably the TF blocking compound binds to tissue factor to thereby inhibit formation of the functional complex. Inhibition or prevention of formation of such a functional complex can have quite broad application, including for treatment of the above-mentioned diseases or disorders in mammals, particularly humans suffering from or susceptible to such diseases or disorders.

Preferred compounds of the invention generally exhibit good blocking activity in at least one test for detecting and preferably measuring TF-mediated blood clotting. More particular tests are standard *in vitro* assays for measuring activity of a specific blood coagulation factor in which the assay is recognized as providing optimal results in the presence of TF or a TF-associated complex such as the human TF/VIIa complex. The TF can be provided in the assay as a recombinant molecule or molecule purified from natural sources depending usually on the specific assay selected.

A more particular *in vitro* assay detects and measures activity of a specific blood coagulation factor which has a recognized activity enhanced in the presence of human TF or the human TF/VIIa complex. Of preferred interest are standard *in vitro* assays for measuring TF-dependent activation of factor X to FXa and factor IX to FIXa. Sometimes these assays will be referred to herein as a "primary screening assay" or related term or phrase such as "method of discovery" to denote preferred use of the assay in screening compounds.

For example, a particularly preferred compound of the invention will exhibit good blocking activity in the primary screening assay for measuring TF-dependent activation of factor X to FXa. Additionally preferred compounds will exhibit good blocking activity in the primary screening assay for measuring TF-dependent activation of factor IX to FIXa.

It will be appreciated that by the phrase "good blocking activity" or related phrase is meant preferred use of a compound of this invention to reduce or inhibit TF/VIIa-dependent activation of factor X to FXa and/or factor IX to FIXa. A preferred compound is a synthetic or semi-synthetic compound such as those small molecule compounds disclosed below. More particular disclosure relating to the primary screening assays is provided as follows.

Preferred compounds of this invention will exhibit an  $IC_{50}$  (concentration required to inhibit factor X activation by about 50% relative to a suitable control) of about 100  $\mu$ M or less and preferably about 10  $\mu$ M or less. Additionally preferred compounds will exhibit equivalent or greater than about 70% inhibition of TF- or TF/VIIa dependent FX activation in the assay. In a preferred embodiment, the primary screening assay includes all of the following steps:

- 1) admixing in a suitable assay solution TF/VIIa complex and factor X under conditions conducive to forming factor Xa,
- 2) contacting the solution with a detectably-labeled factor Xa substrate; and
- 3) detecting labeled product in the solution as being indicative of the factor X activation.

Preferred use of this primary screening assay effectively measures the capability of a candidate compound to decrease or eliminate TF- or TF/VIIa dependent factor X activation. The assay is generally flexible and can be manipulated as necessary to test a compound for capability to block factor X activation. For example, the candidate compound can be added at any one or more of

the steps shown above with addition of the compound at step 1) being preferred for many screening applications.

A preferred TF/VIIa complex for use in the method includes TF which has been exposed to conditions conducive to exposing good TF blocking sites. More specific conditions for isolating and using the TF are provided below.

As mentioned above, another primary screening assay is a standard *in vitro* assay for measuring factor IX activation by TF or TF/VIIa. In this example, a preferred compound will exhibit an  $IC_{50}$  (concentration required to inhibit factor IX activation in the assay by about 50% relative to a suitable control) in the assay of about 200  $\mu$ M or less, and preferably about 10  $\mu$ M or less. In a preferred embodiment, the standard assay for measuring the factor IX activation includes all of the following steps:

- 1) admixing in a suitable assay solution TF/VIIa complex with factor IX under conditions conducive to forming factor FIXa,
- 2) contacting the solution with FX and detectably-labeled FXa substrate; and
- 3) detecting labeled product in the solution as being indicative of the factor IXa activation by TF/VIIa.

In preferred embodiments, this screening assay effectively measures capacity or capability of the candidate compound to decrease or eliminate factor IX activation.

The assay is generally sensitive to TF- or TF/VIIa-dependent formation of FIXa and can be used in several ways to test a desired compound for capacity or capability to block the factor IX activation. For example, a compound to be further tested can be added at one or more of the steps shown above with addition of the compound at step 1) being preferred for most screening applications. Typically preferred compounds of this invention will exhibit good blocking activity in this example of the primary screening assay.

A further preferred primary screen of the invention is the Prothrombin Time (PT) test or assay which measures extrinsic pathway clotting. This test is standard in

the field and is routinely used to measure clotting in biological samples such as blood plasma.

More particularly preferred compounds of this invention will exhibit good inhibitory activity in the PT assay. A typically preferred compound will increase plasma clotting time in the PT assay relative to a suitable control by at least about 5% to about 10% (seconds). Preferred use of the PT assay measures TF-mediated blood plasma clot time and is performed as follows:

- 1) providing citrated plasma in a suitable assay solution under conditions conducive to plasma coagulation,
- 2) admixing a suitable tissue factor preparation and calcium in the solution under conditions suitable for initiating plasma clotting; and
- 3) measuring the clot time in the solution to determine the prothrombin clot time (PT).

Preferred use of the PT assay measures capability of the compound tested to prolong the prothrombin clot time. The PT assay is well known in this field and can be employed in one or a combination of ways to test the compound for capacity or capability to increase or block the prothrombin clot time.

Especially preferred compounds of this invention exhibit good activity in at least one of the primary assays mentioned above (factor X, factor IX activation and/or PT tests).

Good inhibition of the TF- or TF/VIIa-dependent activation in any one or more of the above primary screening assays at least in many cases can be attributed to effects of the compound on TF/VIIa and/or FXa activities. As discussed, preferred compounds of the invention are TF-antagonists and generally exhibit good blocking activity in preferred *in vitro* assays for measuring TF-mediated blood coagulation.

Thus it will usually be desirable to further test compounds giving good blocking activity in one or more of the above primary screening assays and in at least one and preferably more than one of the "secondary screening assays" discussed below. Such

secondary assays can facilitate further identification and selection of candidate compounds having desired TF-antagonist activity, *e.g.*, by eliminating from consideration compounds having activity other than desired activity such as compounds impacting protease activity.

5

A variety of secondary assays can be conducted in accord with this invention to further evaluate compounds identified in a primary assay, *e.g.* to further evaluate activity identified in a primary assay or to determine the presence of a certain undesired activity. For example, additionally preferred compounds of this invention will exhibit substantially reduced or negligible activity in other secondary screening assays which are not optimized to measure TF-antagonism. That is, these secondary assays may not be TF dependent. Particular examples of such assays include those formatted to measure thrombin, trypsin, or activated factors such as FXa, FIXa, or FVIIa. Also, preferred compounds exhibit negligible activity in an Activated Partial Thromoplastin Time (APTT) test or assay. More specific examples of such secondary screening assays are provided in the discussion and Examples which follow.

10

15

In any one or all of the assays disclosed herein including the primary screens and secondary tests discussed above, the candidate compound can be provided in the assay as the sole active agent or it can be combined with other agents to be tested including other compounds or compositions of this invention. In this embodiment, the screening assays are particularly useful for detecting and preferably quantifying synergism between the compounds, agents or compositions.

20

A variety of inhibitors against human tissue factor are disclosed herein. These compounds can be used in the screening assays described herein as well as the treatment and prevention methods of this invention.

25


For example, disclosed herein are phosphonate compounds that are sometimes referenced herein as "TF antagonists", "TF blocking compounds" or similar phrase. Preferred compounds of the invention are small molecules and do not include peptide linkage groups. More particular compounds consist of a phosphonate group and a

30



"headpiece". Typically, the headpiece is covalently bound to the phosphonate group and will include or consist of an amine group or a cyclic ring such as an aromatic group. In embodiments in which the headpiece includes the aromatic group, the headpiece will preferably be linked to a phosphonate (preferably bisphosphonate) group by a nitrogen or oxygen atom. Particular aromatic groups are phenyl groups which can be substituted with one or more other groups as discussed below. In embodiments in which the headpiece is an amine group, it will be appreciated that the compound will be representative of a primary or further substituted amine compound.

10 More specifically, preferred compounds of the invention include those of the following Formula I:



I

15 Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O, S, S(O) or S(O<sub>2</sub>);

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably having 1 to about 12 carbons, more preferably 1 to about 6 carbons; optionally substituted alkenyl preferably having from about 2 to 12 carbon atoms, more preferably about 2 to 6 carbons; optionally substituted alkynyl preferably having from about 2 to 12 carbon atoms, more preferably about 2 to 6 carbon atoms; optionally substituted alkoxy preferably having 1 to about 12 carbon atoms, more preferably 1 to about 6 carbon atoms; optionally substituted alkylthio preferably having from about 1 to 12 carbon atoms, more preferably about 1 to 6 carbon atoms; optionally substituted alkylsulfinyl preferably having from about 1 to 12 carbon atoms, more preferably about 1 to 6 carbon atoms; optionally substituted alkylsulfonyl preferably having from about 1 to 12 carbon atoms, more preferably about 1 to 6 carbon atoms; or optionally substituted alkylamino preferably having from about 1 to 12 carbon atoms, more preferably about 1 to 6 carbon atoms;


m is an integer of from 0 (where the hetero atom is directly substituted on the aryl group) to 4, and preferably is 0, 1 or 2;

n is an integer of from 0 to 4, and preferably n is 1 or 2;

p is 1 (where the compound is a bisphosphonate) or 2 (where the compound has a single terminal PO<sub>3</sub> group); and pharmaceutically acceptable salts thereof.

It is understood that in Formula I above, and elsewhere the designation of “(Het)<sub>0 or 1</sub>” specifies that the Het group may be absent (*i.e.* where the Het subscript is zero) or present in a single occurrence (*i.e.* where the Het subscript is one).

Additional preferred compounds include those of the above formula where Ar is a carbocyclic aryl group, particularly phenyl, such as compounds of the following Formula II:



## II


wherein X, Y, Het, X', Y', Z, m, n, and p are the same as defined in Formula I above;

wherein each R<sup>1</sup> is independently halogen (F, Cl, Br, I); amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl preferably having 1 to about 20 carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted alkenyl preferably having 2 to about 20 carbon atoms, more preferably 2 to about 10 carbon atoms, still more preferably 2 to about 6 carbon atoms; optionally substituted alkynyl preferably having 2 to about 20 carbon atoms, more preferably 2 to about 10 carbon atoms, still more preferably 2 to about 6 carbon atoms; optionally substituted alkoxy preferably having 1 to about 20 carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted alkylthio preferably having 1 to about 20


carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted alkylsulfinyl preferably having 1 to about 20 carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted alkylsulfonyl preferably having 1 to about 20 carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted alkylamino preferably having 1 to about 20 carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted alkanoyl preferably having 1 to about 20 carbon atoms, more preferably 1 to about 10 carbon atoms, still more preferably 1 to about 6 carbon atoms; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

q is an integer of from 0 (where the phenyl ring positions are fully hydrogen substituted) to 5, and preferably m is 0, 1 2 or 3; and pharmaceutically acceptable salts thereof.

Of the compounds of the above Formulae I and II, additional compounds include those where the group Het is optionally substituted nitrogen or oxygen, such as compounds of the following Formulae III and IV:




**III**




**IV**

wherein in each of Formula III and IV,  $R^1$ , X, Y, X', Y', Z, q, m, n, and p are the same as defined in Formulae I and II above; and W is hydrogen, optionally substituted alkyl, preferably having 1 to about 8 carbon atoms, more preferably 1 to about 6 carbon atoms; optionally substituted alkenyl, preferably having 2 to about 8 carbon atoms, more preferably 2 to about 6 carbon atoms; optionally substituted alkynyl, preferably having 2 to about 8 carbon atoms, more preferably 2 to about 6 carbon atoms; optionally substituted alkoxy, preferably having 1 to about 8 carbon atoms, more preferably 1 to about 6 carbon atoms; optionally substituted alkylthio, preferably having 1 to about 8 carbon atoms, more preferably 1 to about 6 carbon atoms; optionally substituted alkylsulfinyl, preferably having 1 to about 8 carbon atoms, more preferably 1 to about 6 carbon atoms; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl, preferably having 1 to about 8 carbon atoms, more preferably 1 to about 6 carbon atoms; optionally substituted carbocyclic aryl; or optionally substituted aralkyl; and pharmaceutically acceptable salts thereof.

Additional compounds of Formula III include those where the nitrogen group is a direct (no interposed carbon or other atoms) phenyl ring substituent, and particularly preferred compounds of Formula IV include those where the oxygen is a direct ring substituent or a single methylene group is present, such as compounds of the following Formulae IIIa and IVa:



**IIIa**



**IVa**

wherein  $R^1$ ,  $X'$ ,  $Y'$ ,  $n$  and  $q$  are the same as defined in Formulae I and II above; and pharmaceutically acceptable salts of those compounds.

5 Additional compounds of the invention bind tissue factor (TF) so that FX does not effectively bind to the TF/factor VIIa complex whereby FX is not effectively converted to its activated form (FXa). Preferred compounds of the invention can inhibit TF function by effectively blocking FX binding or access to TF molecules. See, for instance, the results of Example 2 which follows. As used herein, references  
10 herein to "compounds of the invention" are inclusive of compounds of Formulae I, II, III, IIA, IV and IVA above.

In preferred aspects, the invention provides methods for inhibiting blood coagulation and blood clot formation in a mammal, methods for inhibiting thrombin  
15 generation in a mammal, and methods for treating or preventing thromboembolic disorders in a mammal. The methods of the invention in general comprise administering to a mammal, such as a primate particularly a human, a therapeutically effective amount of a compound of the invention.

20 Compounds of the invention are particularly useful to alleviate various diseases impacted by tissue factor (TF). By the term "impacted" is meant that the severity or duration of the disease is increased by presence of the TF according to the recognized assays or tests. Particular diseases include thromboses, especially to prevent or inhibit restenosis, or other thromboses following an invasive medical  
25 procedure such as arterial or cardiac surgery (*e.g.*, angioplasty or endarterectomy), including for prophylaxis of deep vein thrombosis associated with orthopedic or other surgery. Compounds of the invention also can be employed to reduce or even effectively eliminate blood coagulation arising from use of medical implementation (*e.g.*, a catheter, stent, arteriovenous shunt or other medical device). Compounds of  
30 the invention also will be useful for prophylaxis for long term risk for myocardial infarction. Compounds of the invention also will be useful for treatment of thrombotic conditions that may be associated with acute promyelocytic leukemia,

diabetes, multiple myelomas, disseminated intravascular coagulation associated with septic shock, purpura fulminans associated infection, adult respiratory distress syndrome, unstable angina, and thrombotic complications associated with aortic valve or vascular prosthesis.

5

Additional uses for the present compounds include use in the treatment of atherosclerosis, inflammation, and as an anti-angiogenic agent, especially to treat cancers, particularly solid cancers such as cancers residing in the lung, breast, liver, brain or other tissue.

10

Compounds of the invention also can be employed as an anti-coagulant in extracorporeal circulation of a mammal, particularly a human subject. In such methods, one or more compounds of the invention is administered to the mammal in an amount sufficient to inhibit blood coagulation prior to or during extracorporeal circulation such as may be occur with cardiopulmonary bypass surgery, organ transplant surgery or other prolonged surgeries.

15

Compounds of the invention also can be employed in *in vivo* diagnostic methods including *in vivo* diagnostic imaging of a patient.

20

Compounds of the invention also can be used in *in vitro* assays, e.g. to selectively inhibit factor X activation. Such assays of the invention will be useful to determine the presence or likelihood of a patient having blood coagulation or a blood clot.

25

Pharmaceutical compositions also are provided comprising an effective amount of one or more compounds of the invention and a pharmaceutically acceptable carrier.

30

Other aspects of the invention are discussed *infra*.

## BREIF DESCRIPTION OF THE DRAWINGS

Figure 1 is a flow chart of a screening assay employed in examples below.

Figures 2(a), 2(b), 2 (c) and 2(d) disclose results of examples which follow.

5

Figure 3 is a table showing  $IC_{50}$  values and inhibition of protease activities by certain TF antagonists of this invention, *i.e.* compounds 1, 2, 3, 4, and 6. The compound labeled "Fosamax" is represented by the formula  $NH_2$ -bisphosphonate.

10

Figure 4 is a table showing the effect of specific TF antagonists (compounds 1 and 2) in the prothrombin time (PT) assay.

Figure 5 is a table showing effects of compound 1 on  $K_m$  values for FX in TF/VIIa- dependent activation assay.

15

Figure 6 is a graph showing inhibition of TF/VIIa-dependent activation by compound 1. FX concentration was 30 nM in the assay.

20

Figure 7 is a graph showing inhibition of TF/VIIa-dependent FX activation by compound 1. FX concentration was 5 nM in the assay.

## DETAILED DESCRIPTION OF THE INVENTION

As discussed, the present invention features compounds such as pharmaceutically active compounds and especially pharmaceutical compositions that utilize or comprise one or more of such compounds. Preferred compounds are effective TF antagonists as determined by standard *in vitro* screening assays disclosed herein. Especially preferred compounds are very useful for the treatment or prophylaxis of undesired thrombosis. The invention has a wide spectrum of applications including use in screening candidate compounds having significant TF-antagonistic activity.

25

30

As discussed, preferred compounds and compositions of this invention are good TF antagonists that exhibit significant blocking activity in at least one of and preferably all of the primary screening assays (TF- or TF/VIIa-dependent activation of factor X or factor IX, and PT assay). Especially preferred compounds do not exhibit significant blocking activity in the Activated Partial Thromoplastin Time (APTT) assay discussed previously. Further preferred are those compounds of this invention showing insignificant activity in other secondary assays such as those for measuring trypsin, thrombin, factor Xa, factor IXa, and factor VIIa activity as discussed below.

The standard *in vitro* assays disclosed herein are well-known in the field and are generally flexible. Moreover, the assays can be conveniently manipulated to detect and quantify TF-antagonistic activity as needed. The assays are typically compatible with testing compounds or compositions of this invention in the presence of other therapeutic or experimental agents giving good anti-platelet, anti-thrombolytic, or anti-coagulant activity. In addition, the assays can be used to test effects with recognized anti-TF antibodies. In these embodiments, the standard *in vitro* assays are especially useful for detecting and preferably measuring significant co-operative or synergistic effects exhibited by the compounds or compositions of this invention.

A more particular example of the primary screening assay discussed previously is as follows. The assay is standard for measuring TF/VIIa-dependent factor X activation. A preferred compound exhibits an  $IC_{50}$  in the assay of less than about 100 $\mu$ M and preferably less than about 10  $\mu$ M exemplifying good blocking activity in this assay. In a more preferred embodiment, the primary screening preferably includes the following steps.

1) admixing in a suitable assay solution about 0.1nM of human recombinant TF/VIIa complex (lipidated), about 180nM human FX, and between from about 0.5 $\mu$ l to about 10 $\mu$ l of at least one compound to be tested (optionally dissolved in an appropriate vehicle such as water or dimethylsulfoxide (DMSO)) and incubating the reaction at 37°C for a few minutes up to about an hour or more,



- 2) contacting the solution with a suitable chelating agent such as ethylenediaminetetra acetic acid (EDTA) to reduce or stop factor X activation,
- 3) contacting the solution with a detectable amount of a chromogenic substrate specific for FXa (e.g., Spectrozyme FXa or S-2765) and incubating same at 37°C; and
- 4) detecting chromophore produced in the solution as being indicative of the factor X activation.

Reference herein to a "standard assay for measuring TF/VIIa-dependent factor X activation" or similar phrase will preferably refer to the above steps 1)-4). More specific disclosure relating to the assay can be found in Example 2 below in which the standard assay for measuring TF/VIIa- dependent factor X activation is specifically adapted for spectrophotometric detection of FXa produced chromophores at 405nm.

A preferred TF/VIIa complex for use in the method includes TF that has been exposed to conditions suitable for exhibiting good TF blocking sites. Such TF molecules can be obtained by one or a combination of approaches. In one method, human TF is obtained from an overproducing immortalized cell line or an acetone powder derived from human brain. TF is preferably isolated in the presence of at least one non-ionic detergent such as TRITON® X-100 (polyoxyethylene (10) isooctylphenyl ether) under moderate conditions of salt and pH, e.g., 100 mM NaCl and pH 8.0. Preferred amounts of the non-ionic detergent will vary depending on intended use but will generally be in an amount of from between about 0.05% to about 0.5% (w/v). See the General Comments of the examples below for more specific information about isolating human TF.

Additionally preferred TF is exposed to conditions in the standard assay for measuring TF/VIIa-dependent factor X activation. See Example 2 below for more specific disclosure about that standard assay.

Additionally preferred compounds of this invention exhibit good blocking activity in the other primary screening assay for measuring TF/VIIa-dependent factor

IX activation. Preferred compounds exhibit an  $IC_{50}$  in the assay of less than about 200  $\mu M$  with preferably less than about 10  $\mu M$  exemplifying good blocking activity in this assay. In a more particular embodiment, the standard assay preferably includes the following steps:

- 1) admixing in a suitable assay solution about 0.7 nM TF/VIIa complex with 300 nM factor IX and 1000 nM factor X, and from between about 0.5  $\mu l$  to about 10  $\mu l$  of at least one compound to be tested (optionally dissolved in an appropriate vehicle such as water or dimethylsulfoxide (DMSO)) and incubating the solution at 37°C from between about a few minutes up to about an hour under conditions suitable for forming FIXa and FXa;
- 2) contacting the solution with a suitable chelating agent such as EDTA to stop FIX activation;
- 3) contacting the solution with a chromogenic substrate specific for the FXa (e.g., Spectrozyme FXa) and incubating same at 37°C; and
- 4) detecting chromophore in the solution as being indicative of the factor IX activation.

Reference herein to a "standard assay for measuring TF/VIIa dependent factor IX activation" or similar term or phrase will specifically refer to the above steps 1)-4).

See Example 2 below for a more specific illustration of the standard assay adapted for spectrophotometric detection of preferred chromophore at 405nm.

The table in Figure 3 below shows specific  $IC_{50}$  values for specific TF antagonists of the invention, i.e. compound 1, compound 2, compound 3, compound 4, and compound 6, as well as Fosamax. The values were determined in the standard assays for measuring TF/VIIa-dependent factor X activation and TF/VIIa-dependent factor IX activation. As can be seen from the table in Figure 3, these compounds give good blocking activity in these assays.

As discussed, additionally preferred compounds of this invention exhibit good clot time inhibition in the PT assay, preferably an increase in clotting time from between about 20% to at least 100%, and more preferably from between about 20% to

at least 500% relative to a suitable control. Clot times are generally measured in seconds. Preferred PT assays are typically performed by adding a suitable amount (*e.g.* about 1 to 3 nM) of lipidated tissue factor to an assay solution that includes conventionally citrated plasma. The PT assay measures TF-mediated blood plasma  
5 clot time and is preferably performed by conducting the following steps:

- 1) providing about 0.1 ml of citrated human plasma in a suitable assay solution, and combining same with between from about 0.5 $\mu$ l to 10  $\mu$ l of at least one compound to be tested (optionally dissolved in vehicle such as water or dimethylsulfoxide (DMSO)) and incubating same at room temperature for  
10 about 3 to 10 minutes,
- 2) admixing into the solution from between about 0.2 ml (ca. 1-3 nM recombinant human tissue factor) and about 5-10 mM of calcium to initiate plasma clotting; and
- 3) measuring the plasma clot time to determine the prothrombin clot time  
15 (PT).

Reference herein to a "standard PT assay " or similar phrase or term will specifically refer to the above steps 1)-3). See also *Williams Hematology*, 5<sup>th</sup> Ed. (Beutler, E. et al. Eds.) McGraw-Hill, Inc. Health Professions Div., New York, for  
20 more specific disclosure relating to conducting the PT assay.

As mentioned, the present invention provides a variety of assays for detecting and preferably measuring capability of preferred compounds of this invention to antagonize good TF activity. As has also been discussed, certain standard *in vitro*  
25 screening assays are sometimes referred to herein as "secondary screening assays" to denote preferred use with one or more or all of the primary screening assays mentioned previously. Practice of such particular secondary screening assays in conjunction with one or more of the primary screening assays will provide a wide spectrum of useful compounds featuring good anti-TF activity.


30 Secondary screening assays are disclosed herein and include those optimized to detect and preferably measure the catalytic activities of factor VIIa (FVIIa), factor

IXa (FIXa), factor Xa (FXa), thrombin, trypsin, or Russell's viper venom (RVV). In most instances, optimal practice of these assays does not require added TF. Preferred compounds of this invention are specific TF-antagonists and will generally exhibit substantially reduced or negligible activity in these assays. Practice of the secondary screening tests in conjunction with the primary and preferred secondary screening assays discussed previously will facilitate selection of preferred compounds exhibiting highly specific anti-TF activity. Reference herein to "reduced" or "negligible" activity with respect to these secondary screening assays is meant to denote between from about 2% to about 10% of the activity exhibited by a suitable control such as water or DMSO.

As discussed above, the invention provides a wide spectrum of pharmaceutically active compounds and compositions that are useful to treat or prevent undesired thrombosis. Preferred compounds are tissue factor (TF) antagonists and preferably can specifically block human factor X and IX activation catalyzed by human tissue factor/factor VIIa complex. Illustrative compounds of the invention include the anti-coagulant phosphonate of the above-defined Formula I, II, III, IIIA, IV, and IVA.

Illustrative compounds of the invention are bisphosphonate, *i.e.* compounds of the above formulae where p is 1 and two  $-PO_3$  groups are present. Preferred  $R^1$  ring substituents of the above formulae include hydroxy, halogen, alkyl such as  $C_{1-6}$  alkyl, amino, and alkylamino such as mono-or di- $(C_{1-4})$ alkyl. Preferred W groups (optional amino substituent) include hydrogen, and optionally substituted alkyl, particularly  $C_{1-6}$  optionally substituted alkyl. Preferred X, Y, X', Y' and Z groups include hydrogen and optionally substituted alkyl, particularly  $C_{1-6}$  optionally substituted alkyl.

Additional compounds of the invention include the following compounds 1 through 6, and pharmaceutically acceptable salts of those compounds. Compounds 1, 2 and 6 below are particularly preferred. Those compound designations 1 through 6 are used throughout the present disclosure and refer to the specified compounds of the structures shown immediately below.



Suitable halogen substituent groups of compounds of the invention (which includes e.g. compounds of Formulae I, II, III, IIIA, IV and/or IVA as those formulae are defined above) are F, Cl, Br and I. As used herein, the term alkyl unless otherwise modified refers to both cyclic and noncyclic groups, although cyclic groups will comprise at least three carbon ring atoms. Alicyclic alkyl groups are generally preferred. Alkenyl and alkynyl groups of compounds of the invention have one or more unsaturated linkages, typically 1 to about 3 or 4 unsaturated linkages. Also, the terms alkenyl and alkynyl as used herein refer to both cyclic and noncyclic groups, although straight or branched noncyclic groups are generally more preferred. Alkoxy groups of compounds of the invention have one or more oxygen linkages, typically 1 to about 5 or 6 oxygen linkages. Alkylthio groups of compounds of the invention have one or more thioether linkages, typically 1 to about 5 or 6 thioether linkages. Alkylsulfinyl groups of compound of the invention have one or more sulfinyl (SO) linkages, typically 1 to about 5 or 6 sulfinyl linkages. Alkylsulfonyl groups of compounds of the invention have one or more sulfonyl (SO<sub>2</sub>) linkages, typically 1 to about 5 or 6 sulfonyl linkages. Preferred alkylamino groups of compounds of the invention include those groups having one or more primary, secondary and/or tertiary amine groups, preferably 1 to about 3 or 4 amine groups. Suitable alkanoyl groups have one or more carbonyl groups, typically 1 to about 4 or 5 carbonyl groups.

Alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl and other groups may be suitably either linear or branched. Carbocyclic aryl as used herein refers to non-hetero aromatic groups that have 1 to 3 separate or fused rings and 6 to about 18 carbon ring members and may include e.g. phenyl, naphthyl, biphenyl, acenaphthyl,

5 phenanthracyl, and the like. Phenyl and naphthyl are often preferred. Suitable heteroaromatic or heteroaryl groups will have 1 to 3 rings, 3 to 8 ring members in each ring and from 1 to about 3 hetero atoms (N, O or S). Specifically suitable heteroaromatic or heteroaryl groups include e.g. coumarinyl, quinolinyl, pyridyl, pyrazinyl, pyrimidinyl, furyl, pyrrolyl, thienyl, thiazolyl, oxazolyl, imidazolyl, indolyl,  
10 benzofuranyl, and benzothiazol.

Compounds of Formulae II, III, IIIA, IV and/or IVA as those formulae are defined above preferably have a R<sup>1</sup> group present as a para substituent on the phenyl ring.

15

As discussed above, R', W, X, Y, X', Y', nitrogen "Het" groups, and Z groups are optionally substituted. Suitable groups that may be present on a "substituted" R', W, X, Y, X', Y', Het and Z substituent include e.g. halogen such as fluoro, chloro, bromo and iodo; cyano; hydroxyl; nitro; azido; sulfhydryl; alkanoyl e.g. C<sub>1-6</sub> alkanoyl  
20 group such as acetyl and the like; carboxamido; alkyl groups including those groups having 1 to about 12 carbon atoms, preferably from 1 to about 6 carbon atoms; alkenyl and alkynyl groups including groups having one or more unsaturated linkages and from 2 to about 12 carbon atoms, preferably from 2 to about 6 carbon atoms; alkoxy groups having one or more oxygen linkages and from 1 to about 12 carbon  
25 atoms, preferably 1 to about 6 carbon atoms; aryloxy such as phenoxy; alkylthio groups including those moieties having one or more thioether linkages and from 1 to about 12 carbon atoms, preferably from 1 to about 6 carbon atoms; alkylsulfinyl groups including those moieties having one or more sulfinyl linkages and from 1 to about 12 carbon atoms, preferably from 1 to about 6 carbon atoms; alkylsulfonyl  
30 groups including those moieties having one or more sulfonyl linkages and from 1 to about 12 carbon atoms, preferably from 1 to about 6 carbon atoms; aminoalkyl groups such as groups having one or more N atoms and from 1 to about 12 carbon atoms,

preferably from 1 to about 6 carbon atoms; carbocyclic aryl having 6 or more carbons, particularly phenyl; aryloxy such as phenoxy; aralkyl having 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms, with benzyl being a preferred group; aralkoxy having 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms, with O-benzyl being a preferred group; or a heteroaromatic or heteroalicyclic group having 1 to 3 separate or fused rings with 3 to about 8 members per ring and one or more N, O or S atoms, e.g. coumarinyl, quinolinyl, pyridyl, pyrazinyl, pyrimidyl, furyl, pyrrolyl, thienyl, thiazolyl, oxazolyl, imidazolyl, indolyl, benzofuranyl, benzothiazolyl, tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, morpholino and pyrrolidinyl. A "substituted" R', W, X, Y and Z substituent of a compound of the invention may be substituted at one or more available positions, typically 1, 2 or 3 positions, by one or more suitable groups such as those listed immediately above.

Compounds of the invention can be prepared by procedures generally known in the art. For example, phosphonate acids of Formula I can be prepared by formation of the corresponding alkyl diester followed by conversion to the diacid, e.g. by treatment of the diester with bromotrimethylsilane, and then further reaction of that intermediate to provide a compound of the invention. See, for instance, C. R. Degenhardt et al., *J. Org. Chem.*, 51:3488-3490 (1986); I. S. Alfer et al., *Izv. Akad. Nauk SSSR*, 1122-1126 (1984); I. S. Alfer et al., *Izv. Akad. Nauk SSSR*, 2802-2806 (1983); and U.S. Patent 5,728,650. See also Example 1 which follows.

Reference herein to a "TF blocking compound," "TF antagonist" or related term generally includes those compounds disclosed herein exhibiting good blocking activity in at least one of the primary screening assays such as the PT assay. More particular TF blocking compounds specifically bind TF. Without wishing to be bound to theory, the compounds are believed to block FX or FIX from binding TF in a way sufficient to reduce or block activation to FX or FIX, respectively.

Reference to a "therapeutically effective amount" of a composition is such as to produce a desired effect in a host such as a mammal and especially a primate such

as a human patient. Preferably the effect can be monitored using several end-points known to those of skill in the field. For example, one desired effect is an increase or stabilization of cardiovascular function as measured, e.g., by enhanced heart function and especially blood flow within subject vessels. Such impact can be monitored and usually measured in terms of a therapeutic effect, e.g., improved cardiovascular function, alleviation of one or more symptoms indicative of compromised heart function or function of related vasculature, or other particularized physiological assays. These specific methods are not intended to be inclusive and further methods intended to suit a specific application such as thrombosis, cancer, or atherosclerosis will be apparent to the skilled worker in the field.

As discussed above, a compound of the invention can be administered to a mammal, preferably a primate such as a human, to prevent or reduce thromboses. Therapies in which compounds of the invention will be useful include treatment or prophylaxis of venous thrombosis and pulmonary embolism, arterial thrombosis *e.g.* myocardial ischemia, myocardial infarction, unstable angina, stroke associated with thrombosis, and peripheral arterial thrombosis. Compounds of the invention also may be useful for treatment or prophylaxis of atherosclerotic diseases *e.g.* coronary arterial disease, cerebral arterial disease and peripheral arterial disease. See *e.g.*, Wilde, R. G. et al. *Bioinorganic & Medicinal Chemistry Letters* 167-172 (1995). Compounds of the invention also will be useful for anticoagulation treatment involving artificial organs, cardiac valves, medical implementation (*e.g.* an indwelling device such as a catheter, stent, *etc.*) and the like. Compounds of the invention also will be useful for therapy in other disorders or diseases where blood coagulation may be involved as a related disorder, *e.g.* cancer, inflammatory diseases particularly arthritis, and diabetes.

One or more compounds also may be administered as the sole therapeutic agent(s) in a particular protocol, or the compound(s) of the invention may be administered together with other therapeutic agents, *e.g.* a pharmaceutical targeted for interaction in the blood clotting mechanism such as streptokinase, tPA, urokinase and other agents that lyse clots. A compound of the invention also can be administered with other agents such as one or more other anti-coagulants (*e.g.*, heparin, hirudin, or



hirulog), or an anti-platelet (*e.g.*, ReoPro or aspirin). In such combination therapy, a compound of the invention may be administered prior to, or after administration of one or more other suitable anti-coagulant, anti-platelet, thrombolytic or other agents to boost or prolong desired anti-coagulation activity.

5

Compounds of this invention can be administered intranasally, orally or by injection, *e.g.*, intramuscular, intraperitoneal, subcutaneous or intravenous injection, or by transdermal, intraocular or enteral means. Intravenous or parenteral administration includes *e.g.* sub-cutaneous, intraperitoneal or intramuscular administration. Generally preferred is oral administration. The optimal dose can be determined by conventional means. Compounds of the present invention are suitably administered to a subject in the protonated and water-soluble form, *e.g.*, as a pharmaceutically acceptable salt of an organic or inorganic acid, *e.g.*, hydrochloride, sulfate, hemi-sulfate, phosphate, nitrate, acetate, oxalate, citrate, maleate, mesylate, *etc.*

15

Compounds of the invention can be employed, either alone or in combination with one or more other therapeutic agents as discussed above, as a pharmaceutical composition in mixture with conventional excipient, *i.e.*, pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral or intranasal application which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof.

20

Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, *etc.* The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds.

30

For parenteral application, particularly suitable are solutions, preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Ampules are convenient unit dosages.

5

For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially degradable coatings, *e.g.*, by microencapsulation, multiple coatings, *etc.*

10

See, in general, *Remington's Pharmaceutical Sciences*, (Mack Publishing Co., Easton PA, (1980)), for a discussion of suitable administration formulations.

15

It will be appreciated that the actual preferred amounts of active compounds used in a given therapy will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, the particular site of administration, *etc.* Optimal administration for a given protocol of administration can be readily ascertained by those skilled in the art using conventional dosage determination tests conducted with regard to the foregoing guidelines. In general, a suitable effective dose of one or more compounds of Formula I will be in the range of about 0.01 to 100 milligrams per kilogram of bodyweight of recipient per day, preferably in the range of from about 0.01 to 20 to 50 milligrams per kilogram or bodyweight of recipient per day.

20

25

All documents mentioned herein are fully incorporated by reference in their entirety.

30

The following non-limiting examples are illustrative of the invention.

## General Comments

In the examples below, purified human factors VIIa, IX and X, thrombin, and Russell's viper venom were obtained from Enzyme Research Laboratories Inc.

Trypsin was from Boehringer Mannheim. Chromogenic substrates S-2222, S-2288, S-

5 2238, and S-2765 were from DiaPharma Group Inc., and Spectrozyme FXa was from American Diagnostica Inc. Truncated recombinant human tissue factor (e.g. composed of 243 amino acids) is expressed in *E. coli* and purified by immunoaffinity chromatography. A preferred truncated recombinant human tissue factor lacks the cytoplasmic domain. Native human TF was extracted from human carcinoma cell  
10 line J82 with 50 mM Tris-HCl, pH 8.0, containing 0.1 M NaCl, 1 mM EDTA, 0.3% Triton X-100. Native TF from other sources is extracted with the same buffer from animal brain acetone powders. All other reagents were from Sigma.

**Example 1:** Preparation of 1-(bisphosphonate)-2-amino(3-  
15 hydroxyphenyl)ethyl (compound 1 above)

The method of Degenhardt et al., *J. Org. Chem.*, 51: 3488-3490 (1986) can be followed to produce the compound. Briefly, paraformaldehyde (104.2 g, 3.47 mol) and di-ethylamine (50.8 g, 0.69 mol) are combined in 2 liters of methanol and the mixture warmed until clear. The heat is removed and  $\text{CH}_2(\text{PO}_3(\text{CH}_2\text{CH}_3)_2)_2$  (200 g,  
20 0.69 mol) is added. The mixture is refluxed for 24 hours, and then an additional 2 liters of methanol is added, and the solution concentrated under reduced pressure at 35°C. 1 liter of toluene is added to the concentrate, and the resulting solution concentrated, and the toluene addition and concentration repeated. The resulting intermediate is then dissolved in 1 liter of dry toluene, p-toluenesulfonic acid  
25 monohydrate (0.50 g) is added and the mixture is refluxed. Resulting methanol is removed, e.g. via a Dean-Stark trap or molecular sieves. After 14 hours the solution can be concentrated, diluted in chloroform, washed with water (2 x 150 ml), dried over  $\text{MgSO}_4$  and concentrated. The resulting compound,  $\text{CH}_2=\text{C}(\text{PO}_3(\text{CH}_2\text{CH}_3)_2)_2$ , can be purified if desired such as distillation. The compound  $\text{CH}_2=\text{C}(\text{PO}_3(\text{CH}_2\text{CH}_3)_2)_2$   
30 can then be reacted as desired to provide compounds of the invention. In particular, to provide the title compound,  $\text{CH}_2=\text{C}(\text{PO}_3(\text{CH}_2\text{CH}_3)_2)_2$ , can be reacted with  $\text{NH}_2(3\text{-hydroxyphenyl})$  in a Michael reaction. The phosphono di-ester can be converted to

the di-acid by treatment with bromotrimethylsilane (see, e.g. Morita et al., *Bull. Chem. Soc. Jpn.*, 54:267 (1981)).

### **Example 2: Screening**

5           The primary screening for compounds that inhibit tissue factor/factor VIIa (TF/VIIa) is based on TF/VIIa-dependent FX activation assay (See flow chart in Figure 1 of the drawings). In this assay, the ability of TF/VIIa complex to activate FX is determined in two discontinuous stages. In the first stage (FX activation), the inactive FX is converted to an active enzyme form, FXa, by TF/VIIa in the presence  
10           of phospholipids and calcium. In the second stage (FXa activity assay), EDTA is added at indicated times to the FX activation mixture to chelate calcium, thus leading to the termination of FX --> FXa conversion. Calcium is required for TF/VIIa activity. The activity of FXa is then measured by FXa-specific chromogenic substrates such as S-2222, S-2765, or Spectrozyme FXa. In the primary screening, compounds  
15           from a previously prepared chemical library are first tested at relatively high concentrations (~0.833 mM) in TF/VIIa-dependent FX activation assay to identify hits of potential TF pathway antagonists (see figure 1). However, it is evident that the inhibition of TF/VIIa-dependent FX activation by a compound in this enzyme-coupled assay can be attributed to effects of the compound on TF/VIIa and/or FXa  
20           activities. Thus, secondary screening tests are designed to determine how inhibition takes place and the inhibition mechanism. In secondary screening experiments, effects of those compounds identified from primary screening are tested on catalytic activities of factor VIIa (FVIIa), factor IXa (FIXa), factor Xa (FXa), thrombin, Russell's viper venom (RVV), and trypsin. Additional tests such as the TF/VIIa-dependent factor IX  
25           activation assay and the prothrombin time (PT) assay were conducted to confirm desired activity, and secondary tests were conducted to further select compounds with good TF-antagonistic activity and that did not exhibit undesired activity.

#### **A. Primary Screening: TF/VIIa-dependent FX Activation**

30           Primary screening was done in duplicate in 96-well plates using the TF/VIIa-dependent FX activation assay. All compounds to be screened were dissolved in dimethyl sulfoxide (DMSO), other reagents were prepared or diluted in 25 mM

HEPES-NaOH, 5 mM CaCl<sub>2</sub>, 150 mM NaCl, 0.1% BSA, pH 7.5. For assays where TF was used, purified human recombinant TF (100 nM) was first lipidated with phosphatidylcholine (0.07 mg/ml) and phosphatidylserine (0.03 mg/ml) in 50 mM Tris-HCl, pH 7.5, 0.1 % bovine serum albumin (BSA) for 30 minutes at 37°C. A stock solution of TF/VIIa complex was then prepared by combining equal volumes of 100 nM lipidated TF and 100 nM FVIIa. The complex was incubated at 37°C for 30 minutes and then was aliquoted and stored at -70°C for future uses.

For screening assays, 5 µl of each compound (about 10 mM in DMSO) or DMSO were placed in each well of a 96-well plate, followed by adding 45 µl of TF/VIIa complex (0.1 nM). The components in each well were mixed either with pipette tips or by shaking the plate on a Lab-Line titer plate shaker for 30 seconds. After 15 minutes incubation of the plate at a 37°C, 10 µl of human FX (180 nM) was added to each well and mixed as above. The plate was then incubated at 37°C for 3 to 15 minutes, followed by addition of 10 µl of EDTA (400 mM in 144 mM HEPES, 864 mM NaCl, 0.576% BSA, pH adjusted to 7.5) to each well to terminate FX activation. Ten microliters of FXa substrate (5 mM Spectrozyme FXa, or 3.2 mM S-2765) was added to each well to measure FXa activity. The plate was mixed as above, and after about a 15 minute incubation at 37°C, FXa activity was quenched with 20 µl of 50% acetic acid. Absorbance at 405 nm was then read by an ELISA reader. The OD<sub>405nm</sub> values were transferred to a Microsoft Excel file and the percent inhibition of TF/VIIa-dependent FX activation was calculated by the following formula:

$$\% \text{ Inhibition} = 100 - (100 \times A/B)$$

where A and B are the OD values in the presence and absence of a compound, respectively. Any compound showing equivalent or greater than 70% inhibition of TF/VIIa-dependent FX activation was designated as a candidate for secondary screening test.

## B. Secondary Screening

Those compounds identified in primary screening were retested in TF/VIIa-dependent FX activation assay at 10-, 50- or 100-fold diluted concentrations (see flow

chart of Figure 1 of the drawings). Compounds that failed to show significant inhibition at diluted concentrations, indicating that the inhibition is either non-specific or very weak, were not tested further. Compounds that inhibited TF/VIIa-dependent FX activation at diluted concentrations were further tested for their ability to inhibit activities of the following proteases, trypsin, RVV, thrombin, FXa, FVIIa, and FIXa. A target was to identify compounds that specifically prevent FX (and FIX) binding to TF/VIIa complex or interfere with TF and VIIa interaction so that FX (and FIX) activation is blocked. However, those compounds that have broad ability (non-specific) to inhibit several protease activities were not further investigated.

Compounds that met the specified criteria, that is, to inhibit TF/VIIa-dependent FX activation at lower concentrations ( $< 0.1$  mM) but without significant effects on protease activities, were selected, including compounds 1 and 2, whose structures are shown above, identified as strong TF antagonists and investigated further.

### **Example 3: Effects of Compounds of the Invention on FVIIa, FXa, Thrombin, and Trypsin**

To test whether compounds 1 and 2 inhibit coagulation proteases and trypsin, the following assays were conducted.

#### **A. FVIIa Activity Assay**

Factor VIIa (FVIIa) activity, or the effect of TF and FVIIa interactions, can be determined in the presence of TF using FX and a small peptide (chromogenic) substrate or in the absence of TF using FX as substrate. Assays using FVIIa-specific chromogenic substrate S-2288 directly measures the effect of a compound on FVIIa catalytic activity. In this assay, 55  $\mu$ l of TF/FVIIa complex (containing 10 nM TF and 10 nM VIIa) was first incubated with 5  $\mu$ l of DMSO (or diluted DMSO) or compound, in a 96-well plate for 15 minutes at 37°C, then admixed with 20  $\mu$ l of 8 mM S-2288. The reaction was incubated for 1-2 hours at 37°C. Absorbance at 405 nm was then measured after the reaction was quenched with 20  $\mu$ l of 50% acetic acid. The percent inhibition of TF/VIIa activity was calculated from  $OD_{405nm}$  values in the absence and presence of a compound. Results are shown in Table 1 (Figure 2(a)) and

show that compounds 1 and 2 do not have significant effect on TF/VIIa catalytic activity toward S-2288, indicating that these compounds do not bind to the active site of FVIIa, nor do they interfere with TF and VIIa interactions. Inhibition of TF/VIIa activity toward S-2288 by the two compounds would be expected if they were to bind to VIIa active site or prevent TF and VIIa from forming an active complex.

#### B. FXa Activity Assay

FX is converted to FXa by TF/VIIa complex in the absence of any compound. To do that, 54  $\mu$ l of TF/VIIa (50 nM) was added to 27 ml of buffer in a 50-ml tube. Then 6 ml of FX (180 nM) was added and incubated at 37°C for 15 minutes. Six ml of EDTA was added to stop FX activation. Five  $\mu$ l of compound or DMSO was placed in each well of a 96-well plate in duplicate, then 65  $\mu$ l of FXa generated above was added to each well and mixed. After incubation for 15 minutes at 37°C, 10  $\mu$ l of FXa substrate Spectrozyme FXa was added and incubated for 20 minutes at 37°C. Absorbance at 405 nm was then measured following addition of 20  $\mu$ l of 50% acetic acid. The percent inhibition of FXa activity was calculated from OD<sub>405nm</sub> values in the absence and presence of a compound. Results shown in Table 1 (Figure 2(a)) indicate that compounds 1 and 2 do not inhibit FXa activity, suggesting that inhibition of TF/VIIa-dependent FX activation is not due to the inhibition of FXa activity by these two compounds.

#### C. Thrombin Activity Assay

For thrombin activity assay, 55  $\mu$ l of buffer was mixed with 5  $\mu$ l of DMSO or compound, followed by addition of 10  $\mu$ l of thrombin (20 nM). Mix and incubate at 37°C for 10 minutes. Then 10  $\mu$ l of substrate S-2238 was added and allowed to incubate at 37°C for 15-20 minutes. Absorbance at 405 nm was then measured and the percent inhibition of thrombin activity was calculated from OD<sub>405nm</sub> values in the absence and presence of a compound. Results shown Table 1 (Figure 2(a)) indicated that compounds 1 and 2 (structures shown above) do not inhibit thrombin activity.

#### D. Trypsin Activity

For trypsin activity assay, 4  $\mu$ l of trypsin (100 nM) was first mixed with 61  $\mu$ l of buffer, 5  $\mu$ l of DMSO or 5  $\mu$ l of compound (in DMSO), followed by 15 minute incubation at 37°C. Then 10  $\mu$ l of substrate S-2222 (4.8 mM) was added to start the reaction. After a 15 minute incubation at 37°C, 20  $\mu$ l of 50% acetic acid was added to quench the reaction. Absorbance at 405 nm was then measured and the percent inhibition of trypsin activity was calculated from OD<sub>405nm</sub> values in the absence and presence of a compound. Results in Table 1 (Figure 2(a)) showed that the compounds 1 and 2 (structures shown above) do not inhibit trypsin activity.

See also Figure 3 showing percent inhibition of factor Xa and factor VIIa activity using 83  $\mu$ M compound 1 or compound 2. Figure 3 also shows percent inhibition of thrombin activity at 63  $\mu$ M compound 1 or compound 2. Also shown in the figure is percent inhibition of trypsin activity at 71  $\mu$ M compound 1 or compound 2.

#### **Example 4:** Effects of TF Antagonists on FX Activation Catalyzed by RVV, FIXa, and FVIIa

In addition to TF/VIIa complex, RVV, FIXa, and FVIIa are also able to activate FX *in vitro*. The following assays were conducted to examine the effects of 1 and 2 (structures shown above) on FX activation catalyzed by RVV, FIXa, and VIIa. Data from these assays will help understand where these compounds may bind and the inhibitory mechanism.

##### A. RVV-dependent FX Activation

45  $\mu$ l of RVV (0.1 nM) was added into each well of a 96-well plate that contains 5  $\mu$ l of diluted DMSO (or buffer) or compound, then mixed and incubated at 37°C for 15 minutes. Then 10  $\mu$ l of FX (180 nM) was added and incubated for 15 minutes at 37°C. After adding 10  $\mu$ l of EDTA (400 mM) and 10  $\mu$ l of Spectrozyme FXa (5 mM), the reaction was incubated for 20 minutes at 37°C. Absorbance at 405 nm was then measured following addition of 20  $\mu$ l of 50% acetic acid. The percent



inhibition of RVV-dependent FX activation was calculated from OD<sub>405nm</sub> values in the absence and presence of a compound. The data shown in Table 2 (Figure 2(b)) indicated that the compounds 1 and 2 (structures shown above) do not inhibit RVV catalytic activity, and they do not bind to the cleavage site of FX. It has been established independently that all FX-activating enzymes (TF/VIIa, FVIIa, FIXa, and RVV) cleave the same site on FX.

#### B. FIXa-dependent FX Activation assay:

FIX was first converted to FIXa by TF/VIIa in the absence of compound. This was done in a 50-ml tube. After incubating TF/VIIa (0.67 nM, 10 ml) at 37°C for 15 minutes, FIX (300 nM, 0.123 ml) was added and incubated at 37°C for 30 minutes. Then 1.54 ml of EDTA (400 mM) was added to stop the FIX activation. Then 65 µl of the above FIXa sample was transferred into wells of a 96-well plate that contain 5 µl of diluted DMSO (or buffer) or compound (in diluted DMSO or buffer). Ten µl of FX (1000 nM), 10 µl of polylysine (300 nM), and 10 µl of Spectrozyme FXa (5 mM) were added to each well and incubate at 37°C until an OD<sub>405nm</sub> value of ~0.8 was reached. Absorbance at 405 nm was then measured following addition of 20 µl of 50% acetic acid. The percent inhibition of FIXa activity was calculated from OD<sub>405nm</sub> values in the absence and presence of a compound. Again, compounds 1 and 2 do not inhibit FIXa activity, nor do they bind to FX in such a way that FX can not be activated by FIXa (Table 2, Figure 2(b)).

#### C. FVIIa-dependent FX Activation:

FVIIa alone in the presence of phospholipids and calcium is also able to activate FX, although at a very low rate. This experiment was designed to examine whether compounds 1 and 2 inhibit FVIIa-dependent FX activation if TF is omitted. Inhibition of FVIIa-dependent-FX activation means that the two compounds may bind to FVIIa, while no inhibition indicates that the two compounds will not bind to FVIIa or FX. This assay was done at relatively high FVIIa concentrations. Six microliters of compound 1 (1 mM in 10% DMSO) or 10% DMSO was mixed with 40 µl of 25 mM HEPES-NaOH, 5 mM CaCl<sub>2</sub>, 150 mM NaCl, 0.1% BSA, pH 7.5 containing phosphatidylcholine (0.07 mg/ml) and phosphatidylserine (0.03 mg/ml). Then 4 µl of

FVIIa (1.5  $\mu$ M) and 10  $\mu$ l of FX (180 nM) were added and mixed. The mixture was incubated for 1 hour at 37°C. Then 10  $\mu$ l of EDTA (400 mM) was added to stop FVII activity by removing the calcium required for factor VII activity. Then 10  $\mu$ l of FXa substrate S-2765 (3.2 mM) was added to measure the FXa activity generated by FVIIa. After 16 minutes incubation at 37°C, 20  $\mu$ l of 50% acetic acid was added to quench the reaction. Absorbance at 405 nm was read and the percent inhibition of FVIIa-dependent FX activation was calculated from OD<sub>405nm</sub> values in the absence and presence of compound 1. The data shown in Table 2 (Figure 2(b)) indicates that compound 1 does not inhibit FVIIa-dependent FX activation, indicating that it does not bind to FVIIa or to FX.

#### D. Inhibition of TF/VIIa-dependent FX Activation

To determine the inhibition of TF/VIIa-dependent FX activation by compounds 1 and 2 at lower concentrations, compounds in DMSO were diluted with 10 mM HEPES-NaOH, pH 7.5 and the assays were then performed as previously described in the primary screening method. Table 3 (Figure 2 (c)) is the titration results of TF/VIIa-dependent FX activation for compounds 1 and 2. In some experiments, both compounds were dissolved in 0.1 M NaOH, then diluted by water to 16.5 mM NaOH. From the data in Table 3, the compounds 1 and 2 (structures shown above) inhibit TF/VIIa-dependent FX activation, with IC<sub>50</sub> (inhibitor concentrations at which 50% of TF/VIIa-dependent FX activation is inhibited) values of 19.0  $\mu$ M for compound 1 and 9.7  $\mu$ M for 2.

#### E. Inhibition of TF/VIIa-dependent FIX Activation

TF/VIIa is not only able to activate FX, but is also able to convert FIX to FIXa. To examine whether the two antagonists 1 and 2 were able to block FIX activation catalyzed by TF/VIIa. FIX activation experiment was conducted. To each well of a 96-well plate, 5  $\mu$ l of diluted DMSO or compound was added, followed by 45  $\mu$ l of TF/VIIa (0.67 nM). After mixing and incubating for 15 minutes at 37°C, 10  $\mu$ l of FIX (300 nM) was added and incubated for 10 minutes at 37°C. 10  $\mu$ l of EDTA (400 nM in 144 mM HEPES, 864 mM NaCl, 0.576% BSA, pH adjusted to 7.5) was added, followed by addition of 10  $\mu$ l of FX (1000 nM), 10  $\mu$ l of polylysine, and 10  $\mu$ l

of Spectrozyme FXa (6 mM). After 3 hour incubation at 37°C, absorbance at 405 nm was measured following addition of 20 µl of 50% acetic acid. The percent inhibition of TF/VIIa-dependent FIX activation was calculated from OD<sub>405nm</sub> values in the absence and presence of a compound. The data in Table 4 (Figure 2(d)) shows that compounds 1 and 2 (structure shown above) inhibit TF/VIIa-dependent FIX activation similarly as seen in TF/VIIa-dependent FX activation.

**Example 5: Inhibition Mechanism of TF Antagonists.**

Examples 3 and 4 above showed that the compounds of the invention are TF specific antagonists. To elucidate the inhibition mechanism of TF antagonists, the following experiments were conducted. Compound 1 was titrated from 0 to 84 µM under two assay conditions using two different FX concentrations. Under one condition, compound 1 was preincubated with TF/VIIa for 15 minutes at 37°C prior to addition of FX. Under another condition, TF/VIIa, FX, and compound 1 were added and mixed simultaneously. One set of experiments was conducted at 5 nM FX, the other set at 30 nM FX. From results shown in Figures 6 and 7, IC<sub>50</sub> values from preincubation experiments are somewhat lower than those from the simultaneous addition experiments (compare 8.3 µM with 18.1 µM when FX was 5 nM, and 12.5 µM with 33.2 µM when FX was 30 nM). Furthermore, IC<sub>50</sub> values increased with increasing FX concentration. For example, IC<sub>50</sub> values increased from 8.3 µM and 18.1 µM to 12.5 µM and 33.2 µM, respectively, when FX increased from 5 nM to 30 nM under the two assay conditions. These data suggest that compound 1 compete each other for binding to TF.

Further kinetic analysis of TF-VIIa-dependent FX activation showed that Compound 1 significantly increased the apparent Km values for FX substrate (see Figure 5). This indicates that (1), compound 1 is a competitive inhibitor for TF/VIIa complex, (2), the binding of compound 1 to TF/VIIa blocks FX binding to TF/VIIa complex. The binding of compound 1 to TF (243 form lipidated or 219 form unlipidated) also was directly observed by isothermal calorimetry analysis.

**Example 6:** Effects of Compounds of the Invention in the Prothrombin Time (PT) Assay.

The prothrombin time (PT) test was conducted as follows:

5

The PT assay was performed at 37°C with an Electra 800 (Medical Automation, Inc.). The PT reaction was initiated by adding 0.2 ml of lipidated recombinant human tissue factor into 0.105 ml of human plasma (Ci-Trol Control Level I, from VWR, Cat. No. 68100-336). 1 ml purified water was added to each vial  
10 of Ci-Trol and mixed to solubilize. If more than one vial was used, it was often helpful to combine them into one container. 5 µl of DMSO or 5 µl of compound was added to each well of the twin-well cuvette that contains 0.1 ml of Ci-Trol. It is helpful to use a pipet with 0.1 ml tip to mix each well. Make sure no air bubbles are in the well. Following mixing the compound (or DMSO) with plasma (Ci-Trol), 0.2  
15 ml of lipidated recombinant tissue factor (1-3 nM) is added to the plasma and clotting times were measured within 10 min. The data in Figure 4 indicate compounds 1 and 2 prolonged TF-initiated PT times significantly.

The invention has been described in detail with reference to preferred  
20 embodiments thereof. However, it will be appreciated that those skilled in the art, upon consideration of the disclosure, may make modification and improvements within the spirit and scope of the invention.

What is claimed is:


1. A method for treating a mammal susceptible to, suffering from, suspected of having or recovering from a disease impacted by tissue factor (TF), the method comprising administering to the mammal a therapeutically effective amount of at least one TF blocking compound to treat the disease.
2. The method of claim 1, wherein the disease is selected from the group consisting of a cardiovascular disease, a blood coagulation disorder, a cell proliferation disorder, post-operative complication, an immune disorder, atherosclerosis, inflammation or cancer.
3. A method of blocking or inhibiting tissue factor-dependent activation of factor X and/or factor IX, comprising contacting tissue factor with a TF blocking compound to thereby inhibit formation of a functional complex of factor X or factor IX with tissue factor or TF VIIA.
4. The method of claim 3 wherein the TF blocking compound binds to tissue factor to thereby inhibit formation of the functional complex.
5. The method of any one of claims 1-4 wherein the tissue factor blocking compound exhibits an  $IC_{50}$  of less than about 100  $\mu M$  in a standard assay for measuring TF/VIIa-dependent factor X activation.
6. The method of any one of claims 1-4 wherein the tissue factor blocking compound exhibits an  $IC_{50}$  of about 200  $\mu M$  or less in a standard assay for measuring TF/VIIa-dependent factor IX activation.
7. The method of any one of claims 1-4 wherein the tissue factor compound comprises at least one phosphonate group.

8. The method of any one of claims 1-4 wherein the tissue factor compound comprises at least one bis-phosphonate group.

9. The method of claim 7 or 8 wherein the tissue factor compound further comprises an optionally substituted carbocyclic aryl group, or an optionally substituted heteroaryl group.

10. The method of claim 7 or 8 wherein the tissue factor compound further comprises an optionally substituted phenyl group.

11. The method of any one of claims 1-4 wherein the TF blocking compound is of the following Formula I:



**I**


wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O, S, S(O) or S(O<sub>2</sub>);

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; and pharmaceutically acceptable salts thereof.

12. The method of any one of claims 1-4 wherein the TF blocking compound is of the following Formula II:



## II

wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O, S, S(O) or S(O<sub>2</sub>);

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

each R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

13. A method for identifying a candidate TF blocking compound, the method comprising screening the compound in at least one primary screening assay using purified tissue factor; and selecting the TF blocking compound.


14. The method of claim 13 further comprising screening the candidate TF blocking compound in at least one secondary screening assay.

15. The method of claim 13 or 14 comprising screening the candidate TF blocking compound in a standard *in vitro* assay for measuring TF/VIIa-dependent factor X activation.
16. The method of claim 13 or 14 comprising screening the candidate TF blocking compound in a standard *in vitro* assay for measuring TF/VIIa dependent factor IX activation.
17. The method of claim 13 wherein the screening assay is a standard prothrombin time (PT) assay.
18. The method of any of claims 13-17 wherein the purified tissue factor is lipidated recombinant human tissue factor.
19. A tissue factor (TF) blocking compound exhibiting an  $IC_{50}$  of less than about 100  $\mu M$  in a standard assay for measuring TF/VIIa-dependent factor X activation.
20. The compound of claim 19 further exhibiting equivalent or greater than about 70% inhibition in the assay.
21. The compound of claim 19 further exhibiting an  $IC_{50}$  of about 200  $\mu M$  or less in a standard assay for measuring TF/VIIa-dependent factor IX activation.
22. The compound of claim 19 further exhibiting at least about a 5% to 20% increase in plasma clotting time relative to a control in a standard prothrombin time (PT) assay.
23. The compound of claim 19 further exhibiting at least 70% inhibition in the assay for measuring TF/VIIa-dependent factor X activation and at least of the following activities: 1) an  $IC_{50}$  of less than about 100  $\mu M$  in a standard assay for measuring TF/VIIa-dependent factor IX activation, and 2) at least about a 5% to 20%



increase in plasma clotting time relative to a control in a standard prothrombin time (PT) assay.

24. The compound of claim 19 wherein the compound is represented by the following Formula I:



**I**


wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O or S;

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; and pharmaceutically acceptable salts thereof.

25. The compound of claim 19 wherein the compound is represented by the following Formula II:



**II**

wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;


Het is optionally substituted N, O, S, S(O) or S(O<sub>2</sub>);

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

each R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

26. The compound of claim 19 wherein the compound is represented by the following Formula III:



### III

wherein each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted

alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;


m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5;

W is hydrogen, optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

27. The compound of claim 19 wherein the compound is represented by the following Formula IIIa:




**IIIa**

wherein R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally

substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl; and

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

28. The compound of claim 19 wherein the compound is represented by the following Formula IV:



#### IV


wherein each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5;

$R^1$  is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

29. The compound of claim 19 wherein the compound is represented by the following Formula IVA:



**IVa**

wherein each  $X'$  and each  $Y'$  is independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

$R^1$  is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

$n$  is an integer of from 0 to 4;  $q$  is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

30. A compound of any one of claims 24 through 27 wherein at least one  $R^1$  is hydroxy and  $m$  is 1 or 2.

31. A pharmaceutical composition comprising a compound of any one of claims 19 to 30 and a pharmaceutically acceptable carrier.

32. A method for treating a mammal suffering from or susceptible to a disease impacted by tissue factor, comprising administering to the mammal an effective amount of a compound or composition of any one of claims 19-31.

33. A method for treating a mammal suffering from or susceptible to a cardiovascular disease, a blood coagulation disorder, a cell proliferation disorder, post-operative complication, an immune disorder, atherosclerosis, inflammation, or cancer, comprising administering to the mammal an effective amount of a compound or composition of any one of claims 19-31.

34. A method of inhibiting blood coagulation in a mammal, comprising administering to the mammal an effective amount of a compound or composition of any one of claims 19 to 31.

35. The method of claim 34 wherein the mammal is suffering from, suspected of having, or recovering from a thrombosis.

36. The method of claim 34 wherein the mammal is suffering from, susceptible to, or recovering from restenosis associated with an invasive medical procedure.

37. The method of claim 34 wherein the invasive medical procedure is angioplasty, endarterectomy, deployment of a stent, use of catheter, graft implantation or use of an arteriovenous shunt.

38. The method of claim 34 wherein the mammal is suffering from, at risk of developing, or recovering from a thromboembolic condition associated with cardiovascular disease, an infectious disease, a neoplastic disease or use of a thrombolytic agent.

39. The method of any one of claims 34-38 wherein at least one of an anti-platelet composition, thrombolytic composition or an anti-coagulant composition is administered in combination with a tissue factor blocking compound exhibiting an  $IC_{50}$  of less than about 100  $\mu$ M in a standard assay for measuring TF/VIIa-dependent factor X activation.


40. A method of treating or preventing a thromboembolic disorder in a mammal, comprising administering to the mammal an effective amount of the TF blocking compound of any one of claims 19-31.

41. The method of claim 40 wherein the mammal is suffering from, at risk developing, or is recovering from a thromboembolic condition associated with cardiovascular disease, an infectious disease, a neoplastic disease or use of a thrombolytic agent or an anti-platelet agent.

42. The method of claim 40 wherein at least one of an anti-platelet, thrombolytic, or an anti-coagulant composition is administered in combination with a tissue factor blocking compound exhibiting an  $IC_{50}$  of less than about 100  $\mu$ M in a standard assay for measuring TF/VIIa-dependent factor X activation.

43. The method of any one of claims 32-42 wherein the mammal is a human.

44. A compound of the following Formula I:



**I**


wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O, S, S(O) or S(O)<sub>2</sub>;

each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; and pharmaceutically acceptable salts thereof.

45. A compound of claim 44 having the following Formula II:



## II

wherein Ar is optionally substituted carbocyclic aryl or optionally substituted heteroaryl;

Het is optionally substituted N, O, S, S(O) or S(O<sub>2</sub>);


each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

each R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.



46. A compound of claim 44 having the following Formula III:



### III

wherein each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;


m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5;

W is hydrogen, optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

$R^1$  is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

47. A compound of claim 44 having the following Formula IIIa:




**IIIa**

wherein R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl; and

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

48. A compound of claim 44 having the following Formula IV:



**IV**


wherein wherein each X, each Y, each X', each Y' and each Z are each independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

m and n each is independently an integer of from 0 to 4; p is 1 or 2; q is an integer of from 0 to 5;

R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl;

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

49. A compound of claim 44 having the following Formula IVa:



**IVa**


wherein each X' and each Y' is independently hydrogen; halogen; hydroxyl; sulfhydryl; amino; optionally substituted alkyl preferably; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; or optionally substituted alkylamino;

R<sup>1</sup> is independently halogen; amino; hydroxy; nitro; carboxy; sulfhydryl; optionally substituted alkyl; optionally substituted alkenyl; optionally substituted alkynyl; optionally substituted alkoxy; optionally substituted alkylthio; optionally substituted alkylsulfinyl; optionally substituted alkylsulfonyl; optionally substituted alkylamino; optionally substituted alkanoyl; optionally substituted carbocyclic aryl; or optionally substituted aralkyl; and

q is an integer of from 0 to 5; and pharmaceutically acceptable salts thereof.

50. A compound of any one of claims 44 through 46 wherein at least one R<sup>1</sup> is hydroxy and m is 1 or 2.

51. A compound of claim 44, that is:



or a pharmaceutically acceptable salt thereof.

52. A method of inhibiting blood coagulation in a mammal, comprising administering to the mammal an effective amount of a compound of any one of claims 44 to 51.

53. The method of claim 52 wherein the mammal is suffering from or suspected of having a thrombosis.

54. The method of claim 52 wherein the mammal is suffering from or susceptible to restenosis associated with an invasive medical procedure.

55. The method of claim 54 wherein the invasive medical procedure is angioplasty, endarterectomy, deployment of a stent, use of catheter, graft implantation or use of an arteriovenous shunt.

56. The method of claim 52 wherein the mammal is suffering from or at risk of developing a thromboembolic condition associated with cardiovascular disease, an infectious disease, a neoplastic disease or use of a thrombolytic agent.

57. The method of any one of claims 52 to 56 wherein an anti-platelet composition, a thrombolytic composition or an anti-coagulant composition is administered in combination with a compound of Formula I.

58. A method of treating or preventing a thromboembolic disorder in a mammal, comprising administering to the mammal an effective amount of a compound of any one of claims 44 to 51.

59. The method of claim 58 wherein the mammal is suffering from or at risk developing a thromboembolic condition associated with cardiovascular disease, an infectious disease, a neoplastic disease or use of a thrombolytic agent or an anti-platelet agent.

60. The method of claim 58 or 59 wherein an anti-platelet composition, a thrombolytic composition or an anti-coagulant composition is administered in combination with a compound of Formula I.

61. A method of treating a mammal suffering from or susceptible to atherosclerosis, comprising administering to the mammal an effective amount of a compound of any one of claims 44 to 51.

62. The method of any one of claims 52 to 61 wherein the mammal is a human.

63. A pharmaceutical composition comprising a compound of any one of claims 44 through 51 and a pharmaceutically acceptable carrier.

#### ABSTRACT OF THE DISCLOSURE

The invention includes pharmaceutically active compounds and methods of treatment and pharmaceutical compositions that utilize or comprise one or more such compounds. Compounds of the invention are particularly useful for treatment or prophylaxis of undesired thrombosis.

#125868

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2

## TF/VIIa-dependent Factor X Activation

TF/VIIa/phospholipids/ $\text{Ca}^{++}$  + DMSO (diluted DMSO or Buffer)

or Compound

15 min at 37 C



Add FX



Add EDTA



Add FXa Substrate



Add 50% Acetic Acid



Read OD at 405nm

Figure 1

**Table 1**

Compounds	% Inhibition of FXa Activity at		% Inhibition of FVIIa Activity at		% Inhibition of Thrombin Activity at		% Inhibition of Trypsin Activity at	
	714 $\mu$ M	71 $\mu$ M	833 $\mu$ M	83 $\mu$ M	625 $\mu$ M	63 $\mu$ M	625 $\mu$ M	63 $\mu$ M
<b>Compound <u>2</u></b>	3	3	27	5	57	-16	-8	-16
<b>Compound <u>1</u></b>	7	1	42	1	41	-9	-2	10

Compounds were in DMSO and were diluted with 10 mM Hepes-NaOH, pH 7.5 if necessary

**Figure 2(a)**



**Table 2**

Compounds	% Inhibition of FX Activation Catalyzed by			
	TF/VII <sub>1</sub>	RVV	FIX <sub>1</sub>	FVII <sub>1</sub>
<b>Compound 2</b>				
96 $\mu$ M	nd*	6.1	23.2	nd
32 $\mu$ M	94.5	1.8	6.8	nd
<b>Compound 1</b>				
100 $\mu$ M	nd	nd	nd	6.0
96 $\mu$ M	nd	5.0	21.7	nd
32 $\mu$ M	85.0	4.9	0.0	nd

nd\*, not determined.

The compounds were in DMSO and were diluted with 10 mM Hepes-NaOH, pH 7.5 if necessary.

**Figure 2(b)**

**Table 3**

Compound Concentrations ( $\mu$ M)	% Inhibition by Compound <u>1</u> <u>2</u>	% Inhibition by Compound <u>1</u>
0	0	0
2	6	1
4	10	7
8	45	14
16	62	42
32	84	83
48	92	91
96	96	94

Compounds 1 and 2 were dissolved in 0.1 M NaOH, then diluted to 16.5 mM NaOH with water. The compounds were preincubated with TF VIIa for 15 minutes at 37 C prior to addition of FX. Following FX addition, the reaction was incubated for 15 minutes at 37 C

**Figure 2(c)**

**Table 4**

Compound Concentrations ( $\mu$ M)	% Inhibition by Compound <u>2</u>	% Inhibition by Compound <u>1</u>
0	0	0
2	24.7	
4	39.9	-0.3
8	67.2	20.8
16	79.4	26.9
32	87.5	61.1
48		84.5
96		90.3

Compounds 1 and 2 were in DMSO and diluted with 10 mM HEPES-NaOH, pH 7.5.

**Figure 2(d)**

Compounds	Factor X Activation IC <sub>50</sub> in $\mu$ M	Factor IX Activation IC <sub>50</sub> in $\mu$ M	FXa Activity % Inhibition at 8 $\mu$ M	FVIIa Activity % Inhibition at 8 $\mu$ M	Thrombin Activity % Inhibition at 6 $\mu$ M	Trypsin Activity % Inhibition at 7 $\mu$ M
Compound 1	10	26	4	1	0	4
Compound 2	6.5	55	3	5	0	0
Compound 3	70	n.d.	n.d.	n.d.	n.d.	n.d.
Compound 4	>200	n.d.	n.d.	n.d.	n.d.	n.d.
Compound 5	100	n.d.	n.d.	n.d.	n.d.	n.d.
Fosamur	80	n.d.	n.d.	n.d.	0	n.d.

n.d. - not determined  
Figure 3

<i>Compounds</i>	<i>PT Clotting Time (Seconds)</i>	<i>APTT Clotting Time (Seconds)</i>
<i>Control (DMSO)</i>	<i>37.0</i>	<i>29.4</i>
<i>Compound 1 (167 <math>\mu</math>M)</i>	<i>75.3</i>	<i>35.4</i>
<i>Compound 2 (167 <math>\mu</math>M)</i>	<i>76.8</i>	<i>33.0</i>

*Figure 4*

Compound	K <sub>m</sub> for FX (nM)
Control (DMSO)	19
10 $\mu$ M - compound 1 (in DMSO)	37
20 $\mu$ M - compound 2 (in DMSO)	76

Figure 5




Figure 6




Figure 7